arXiv:1201.2633v1 [math.NT] 12 Jan 2012

ON THE ASYMPTOTICS OF THE RIEMANN ZETA FUNCTION TO ALL ORDERS

A.S. Fokas$^{(1)}$ and J. Lenells$^{(2)}$

$^{(1)}$ Department of Applied Mathematics and Theoretical Physics,
University of Cambridge, Cambridge, CB3 0WA, UK
T.Fokas@damtp.cam.ac.uk
$^{(2)}$ Department of Mathematics, Baylor University,
One Bear Place #97328, Waco, TX 76798, USA
Jonatan_Lenells@baylor.edu

January 13, 2012

Dedicated to Trevor Stuart with deep gratitude

Abstract

We present several formulae for the large t asymptotics of the Riemann zeta function $\zeta(s)$, $s = \sigma + it$, $0 \leq \sigma \leq 1$, $t > 0$, which are valid to all orders. A particular case of these results coincides with the classical results of Siegel. Using these formulae, we derive explicit representations for the sum $\sum_a^b n^{-s}$ for certain ranges of a and b. In addition, we present precise estimates relating this sum with the sum $\sum_c^d n^{s-1}$ for certain ranges of a, b, c, d. Finally, we derive certain novel integral representations for the basic sum characterising the leading large t asymptotics of $\zeta(s)$.

Contents

1 Introduction **2**
1.1 The large t asymptotics of $\zeta(s)$ valid to all orders 2
1.2 The explicit form of certain sums 5
1.3 The explicit form of the difference of certain sums 5
1.4 Several representations for the basic sum 6

2 An Exact Representation of $\zeta(s)$ **7**

3 The Asymptotics of the Riemann Zeta Function for $t \leq \eta < \infty$ 11

4 The Asymptotics of the Riemann Zeta Function for $0 < \eta < t$ 26

5 Consequences of the Asymptotic Formulae 45

6 Several Representations for the Basic Sum 48

A The Asymptotics of $\Gamma(1-s)$ and $\chi(s)$ 58

B Numerical Verifications 59
- B.1 Verification of Theorem 3.1 59
- B.2 Verification of Theorem 3.2 60
- B.3 Verification of Theorem 4.1 60
- B.4 Verification of Corollary 4.3 61
- B.5 Verification of Theorem 4.4 61

1 Introduction

It is well known, see for example theorem 4.11 in Titchmarsh [9], that the Riemann zeta function $\zeta(s)$, $s = \sigma + it$, $\sigma, t \in \mathbb{R}$, satisfies the equation

$$\zeta(s) = \sum_{n \leq x} \frac{1}{n^s} - \frac{x^{1-s}}{1-s} + O\left(\frac{1}{x^\sigma}\right), \qquad |t| < \frac{2\pi x}{c}, \quad c > 1, \quad \sigma > 0, \quad t \to \infty. \quad (1.1)$$

Here we present a systematic investigation of asymptotic formulae for $\zeta(s)$. In this connection, in section 2 we derive the following exact formula (see lemma 2.1):

$$\zeta(1-s) = \sum_{n=1}^{\left[\frac{\eta}{2\pi}\right]} n^{s-1} + \frac{1}{(2\pi)^s}\left[-\frac{\eta^s}{s} + \left(e^{\frac{i\pi s}{2}} \int_{\eta e^{-\frac{i\pi}{2}}}^{\infty e^{i\phi_1}} + e^{-\frac{i\pi s}{2}} \int_{\eta e^{\frac{i\pi}{2}}}^{\infty e^{i\phi_2}}\right) \frac{z^{s-1}dz}{e^z - 1}\right],$$
$$0 < \eta < \infty, \quad -\frac{\pi}{2} < \phi_j < \frac{\pi}{2}, \quad j = 1, 2, \qquad (1.2)$$

where the contours of integration in the first and second integrals in the rhs of equation (1.2) are the rays from $\eta\exp(-i\pi/2)$ to $\infty\exp(i\phi_1)$ and from $\eta\exp(i\pi/2)$ to $\infty\exp(i\phi_2)$, respectively.

1.1 The large t asymptotics of $\zeta(s)$ valid to all orders

Equation (1.2) suggests a separate analysis for the cases $t < \eta$, $t = \eta$ and $t > \eta$. The first two cases are analysed in theorems 3.1 and 3.2 of section 3. Theorem 3.1 presents the large t asymptotics of $\zeta(s)$ valid to all orders in the case of $(1+\epsilon)t < \eta$.

For example, the formulae of theorem 3.1 yield the following equation for the leading asymptotic terms in the case when $2t < \eta$:

$$\zeta(s) = \sum_{n=1}^{[\frac{\eta}{2\pi}]} n^{-s} - \frac{1}{1-s}\left(\frac{\eta}{2\pi}\right)^{1-s} + \frac{2i\eta^{-s}}{(2\pi)^{1-s}}\left\{-i\arg(1-e^{i\eta}) + \frac{t-i\sigma}{\eta}\operatorname{Re}\operatorname{Li}_2(e^{i\eta})\right\}$$
$$+ O\left(\frac{t^2}{\eta^{2+\sigma}}\right), \qquad 2t < \eta < \infty, \quad 0 \le \sigma \le 1, \quad t \to \infty, \tag{1.3}$$

where the error term is uniform for η, σ in the above ranges and the polylogarithm $\operatorname{Li}_m(z)$ is defined by

$$\operatorname{Li}_m(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^m}, \qquad m \ge 1.$$

Theorem 3.2 presents the large t asymptotics of $\zeta(s)$ valid to all orders in the case of $\eta = t$. The formula of theorem 3.2 yields the following equation for the leading asymptotic terms:

$$\zeta(s) = \sum_{n=1}^{[\frac{t}{2\pi}]} n^{-s} - \frac{1}{1-s}\left(\frac{t}{2\pi}\right)^{1-s}$$
$$- \frac{2it^{-s}e^{it}}{(2\pi)^{1-s}}\left\{-\frac{e^{-it}}{2t}[t + i(\sigma-1)] - i\operatorname{Im}\operatorname{Li}_1(e^{it}) + \frac{i\sigma}{t}\operatorname{Re}\operatorname{Li}_2(e^{it}) - \frac{1}{t}\operatorname{Im}\operatorname{Li}_3(e^{it})\right\}$$
$$+ \frac{t^{-s}e^{it}}{(2\pi)^{1-s}}\left\{\frac{1+i}{2}\sqrt{\pi t} - \frac{1}{3}i(3\sigma-2)\right.$$
$$+ \frac{i-1}{24\sqrt{t}}\sqrt{\pi}\left(6\sigma^2 - 6\sigma + 1\right) + \frac{1}{135t}\left(45\sigma^3 - 45\sigma^2 + 4\right)$$
$$\left. - \frac{1+i}{576t^{\frac{3}{2}}}\sqrt{\pi}\left(36\sigma^4 - 24\sigma^3 - 24\sigma^2 + 12\sigma + 1\right)\right\} + O\big(t^{-\sigma-2}\big),$$
$$0 \le \sigma \le 1, \quad t \to \infty, \tag{1.4}$$

where the error term is uniform for σ in the above range.

The best estimate for the growth of $\zeta(s)$ as $t \to \infty$, is not based on equation (1.1) but on the well known approximate functional equation, see for example equation (4.12.4) on page 79 of Titchmarsh [9]:

$$\zeta(s) = \sum_{n \le x} \frac{1}{n^s} + \chi(s)\sum_{n \le y} \frac{1}{n^{1-s}} + O\left(x^{-\sigma} + |t|^{\frac{1}{2}-\sigma}y^{\sigma-1}\right), \qquad xy = \frac{t}{2\pi}, \quad 0 < \sigma < 1, \tag{1.5}$$

where $\chi(s)$ is defined by

$$\chi(s) = \frac{(2\pi)^s}{\pi}\sin\left(\frac{\pi s}{2}\right)\Gamma(1-s),$$

with $\Gamma(s)$ denoting the Gamma function. Building on Riemann's unpublished notes, Siegel in his classical paper [8] presented the error terms of the rhs of equation (1.5) to *all* orders, only in the important case of $x = y = \sqrt{t/2\pi}$. In theorems 4.1 and 4.4 we present analogous results for *any* x and y valid to *all* orders. Theorem 4.1 yields the following explicit result for the subleading term in the rhs of equation (1.5) when $1 < \eta = 2\pi y < \sqrt{t}$. For every $\epsilon > 0$, there exists an $A > 0$ such that

$$\begin{aligned}\zeta(s) = \sum_{n=1}^{[\frac{t}{\eta}]} \frac{1}{n^s} + \chi(s) \sum_{n=1}^{[\frac{\eta}{2\pi}]} \frac{1}{n^{1-s}} - \frac{e^{-i\pi s}\Gamma(1-s)}{i\sqrt{2\pi t}} e^{-([\frac{t}{\eta}]+1)i\eta} e^{\frac{i\pi}{2}(s-1)} \eta^s \frac{e^{\frac{i\pi}{4}}}{1-e^{-i\eta}} \\ + e^{-i\pi s}\Gamma(1-s) e^{-\frac{\pi t}{2}} \eta^{\sigma-1} \times \begin{cases} O\left(\frac{\eta}{t^{5/6}}\right), & 1 < \eta < At^{\frac{1}{3}} < \infty, \\ O\left(e^{-\frac{At}{\eta^2}} + \frac{\eta^3}{t^{3/2}}\right), & t^{\frac{1}{3}} < \eta < A\sqrt{t} < \infty, \end{cases} \\ \mathrm{dist}(\eta, 2\pi\mathbb{Z}) > \epsilon, \quad 0 \le \sigma \le 1, \quad t \to \infty,\end{aligned}$$

where the error term is uniform for all η, σ in the above ranges. In corollary 4.3, similar formulas are presented for the case $2\pi\sqrt{t} < \eta < \frac{2\pi}{\epsilon} t$.

The case of $\epsilon\sqrt{t} < \eta < t$ is analyzed in theorem 4.4, which provides a formula valid to all orders in terms of the function $\Phi(\tau, u)$ defined by

$$\Phi(\tau, u) = \int_{0 \nwarrow 1} \frac{e^{\pi i \tau x^2 + 2\pi i u x}}{e^{\pi i x} - e^{-\pi i x}} dx, \qquad \tau < 0, \quad u \in \mathbb{C}, \tag{1.6}$$

where $0 \nwarrow 1$ denotes a straight line parallel to $e^{3\pi i/4}$ which crosses the real axis between 0 and 1. The formula of theorem 4.4 yields the following result for the leading asymptotic terms: For every $\epsilon > 0$,

$$\begin{aligned}\zeta(s) = &\sum_{n=1}^{[\frac{t}{\eta}]} \frac{1}{n^s} + \chi(s) \sum_{n=1}^{[\frac{\eta}{2\pi}]} \frac{1}{n^{1-s}} \\ &+ e^{-i\pi s}\Gamma(1-s) \Bigg\{ e^{\frac{i\pi(s-1)}{2}} \eta^{s-1} e^{\frac{2t}{\eta}[\frac{\eta}{2\pi}]\pi i - it - \frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi - \eta)^2} \\ &\times \left[\Phi + \frac{\sigma-1}{i\eta}\left(\partial_2\Phi + \left(2\left[\frac{\eta}{2\pi}\right]\pi i - i\eta\right)\Phi\right)\right] + O\left(e^{-\frac{\pi t}{2}} \frac{\eta^\sigma}{t^{5/6}}\right)\Bigg\}, \\ &\epsilon\sqrt{t} < \eta < t, \quad 0 \le \sigma \le 1, \quad t \to \infty,\end{aligned} \tag{1.7}$$

where the error term is uniform for η, σ in the above ranges and Φ and $\partial_2\Phi$ are evaluated at the point

$$\left(-\frac{2\pi t}{\eta^2}, \frac{2t}{\eta} - \frac{2\pi t}{\eta^2}\left[\frac{\eta}{2\pi}\right] - \left[\frac{t}{\eta}\right] - \frac{1}{2}\right). \tag{1.8}$$

Remark 1.1 1. In the particular case of $\eta = \sqrt{2\pi t}$, equation (1.7) agrees with the analogous formula of Siegel [8].

2. Although Siegel suggested already in [8] the possibility of deriving a formula such as (1.7) based on the function (1.6), the authors have not been able to locate such a formula in the literature. In the case of $\sigma = 1/2$, an alternative very interesting rigorous asymptotic representation for $\zeta(s)$ involving a sum which is smoothly rather than sharply truncated is presented in [3].

3. Equation (1.7) is particularly useful in the case when $\eta = \sqrt{\frac{2\pi t}{b}}$ where $b > 0$ is a rational number. Indeed, in this case the function Φ and its derivatives evaluated at the point (1.8) can be computed explicitly (see equation (4.29) below). Therefore, in this case theorem 4.4 yields an *explicit* asymptotic expansion of $\zeta(s)$ to all orders.

4. Equation (1.3) is only useful as an asymptotic formula in the case when $t/\eta \to 0$ as $t \to \infty$ (otherwise the error term is as large as the retained terms). Therefore the asymptotic range where $t < \eta$ and $t/\eta = O(1)$ is not covered by this equation. However, this asymptotic sector is covered by corollary 4.3.

1.2 The explicit form of certain sums

Theorems 3.1 and 3.2 allow us to evaluate the sum $\sum_a^b n^{-s}$, for certain a and b, to *all* orders, see theorems 5.1 and 5.2. For example, the leading order of such sums follows immediately from equations (1.3) and (1.4): For every $\epsilon > 0$,

$$\sum_{n=\left[\frac{\eta_1}{2\pi}\right]+1}^{\left[\frac{\eta_2}{2\pi}\right]} n^{-s} = \frac{1}{1-s}\left[\left(\frac{\eta_2}{2\pi}\right)^{1-s} - \left(\frac{\eta_1}{2\pi}\right)^{1-s}\right] + O(\eta_1^{-s}), \tag{1.9}$$

$$(1+\epsilon)t < \eta_1 < \eta_2 < \infty, \quad 0 \le \sigma \le 1, \quad t \to \infty,$$

$$\sum_{n=\left[\frac{t}{2\pi}\right]+1}^{\left[\frac{\eta}{2\pi}\right]} n^{-s} = \frac{1}{1-s}\left[\left(\frac{\eta}{2\pi}\right)^{1-s} - \left(\frac{t}{2\pi}\right)^{1-s}\right] + O(t^{-s}), \tag{1.10}$$

$$(1+\epsilon)t < \eta < \infty, \quad 0 \le \sigma \le 1, \quad t \to \infty,$$

uniformly with respect to η_1, η_2, η, and σ.

1.3 The explicit form of the difference of certain sums

It is well known, see for example page 78 of Titchmarsh [9], that for large t, the following sums coincide to the leading order:

$$\sum_{x<n\le N} n^{-s} \qquad \text{and} \qquad \chi(s) \sum_{\frac{t}{2\pi N}<n\le\frac{t}{2\pi x}} n^{s-1}.$$

Theorems 4.1 and 4.4 allow us to evaluate the difference of similar sums to all orders, see theorem 5.3. In particular, equation (1.7) immediately implies the following equation:

$$\sum_{n=[\frac{t}{\eta_2}]+1}^{[\frac{t}{\eta_1}]} \frac{1}{n^s} = \chi(s) \sum_{n=[\frac{\eta_1}{2\pi}]+1}^{[\frac{\eta_2}{2\pi}]} \frac{1}{n^{1-s}} \tag{1.11}$$

$$+ e^{-i\pi s}\Gamma(1-s)\Bigg\{ e^{\frac{i\pi(s-1)}{2}} \eta^{s-1} e^{\frac{2t}{\eta}[\frac{\eta}{2\pi}]\pi i - it - \frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi - \eta)^2}$$

$$\times \left[\Phi + \frac{\sigma-1}{i\eta}\left(\partial_2\Phi + \left(2\left[\frac{\eta}{2\pi}\right]\pi i - i\eta\right)\Phi\right)\right] + O\left(e^{-\frac{\pi t}{2}}\frac{\eta^\sigma}{t^{5/6}}\right)\Bigg\}\Bigg|_{\eta=\eta_1}^{\eta_2},$$

$$\epsilon\sqrt{t} < \eta_1 < \eta_2 < t, \quad 0 \le \sigma \le 1, \quad t \to \infty.$$

1.4 Several representations for the basic sum

The remarkable fact about the large t-asymptotics of the Riemann zeta function is that, whereas the higher order terms in the asymptotic expansion can be computed *explicitly*, the sum appearing in the leading order is *transcendental*. In section 6 we present several integral representations of this basic sum with the hope that these representations may be useful for the estimation of this fundamental sum (some results in this direction can be found in [1]).

Let C_η^t denote the semicircle from $i\eta$ to it with $\operatorname{Re} z \ge 0$:

$$C_\eta^t = \left\{ \frac{i(\eta+t)}{2} + \frac{t-\eta}{2}e^{i\theta} \;\middle|\; -\frac{\pi}{2} < \theta < \frac{\pi}{2} \right\}. \tag{1.12}$$

The following representations are derived in section 6:

$$\begin{aligned} \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s}\int_{C_\eta^t} \frac{z^{s-1}}{e^z-1}dz &= \frac{1}{(2\pi)^s}\lim_{\substack{\epsilon\to 0\\ \epsilon t \to 0}} \int_\eta^t \frac{u^{s-1}}{e^{iue^{-i\epsilon}}-1}du \\ &= \frac{2}{(2\pi)^s} \fint_\eta^t \frac{\rho^{s-1}}{e^{i\rho}-1}d\rho + O(t^{\sigma-1}) \\ &= \sum_{n=[\frac{\eta}{2\pi}]+1}^{[\frac{t}{2\pi}]} n^{s-1} + O(t^{\sigma-1}), \qquad 0 < \eta < t, \quad t \to \infty, \end{aligned}$$

where it is assumed that $\operatorname{dist}(\eta, 2\pi\mathbb{Z}) > \delta$, $\operatorname{dist}(t, 2\pi\mathbb{Z}) > \delta$, and the contour in the second equality above denotes the principal value integral with respect to the points

$$\left\{ 2\pi n \;\middle|\; n \in \mathbb{Z}^+, \ \left[\frac{\eta}{2\pi}\right] + 1 \le n \le \left[\frac{t}{2\pi}\right] \right\}.$$

Basic Assumption: Throughout this paper it is assumed that

$$\eta > 0, \quad t > 0, \quad \eta \notin 2\pi\mathbb{Z}, \quad t \notin 2\pi\mathbb{Z}. \tag{1.13}$$

2 An Exact Representation of $\zeta(s)$

In this section, we assume that the branch cut for the logarithm runs along the negative real axis.

Lemma 2.1 (An exact representation of Riemann's zeta function) *Let* $\zeta(s)$, $s = \sigma + it$, $\sigma, t \in \mathbb{R}$, *denote the Riemann zeta function. Then,*

$$\zeta(s) = \chi(s)\left\{\sum_{n=1}^{\left[\frac{\eta}{2\pi}\right]} n^{s-1} + \frac{1}{(2\pi)^s}\left[-\frac{\eta^s}{s} + \left(e^{\frac{i\pi s}{2}}\int_{\eta e^{-\frac{i\pi}{2}}}^{\infty e^{i\phi_1}} + e^{-\frac{i\pi s}{2}}\int_{\eta e^{\frac{i\pi}{2}}}^{\infty e^{i\phi_2}}\right)\frac{e^{-z}z^{s-1}dz}{1-e^{-z}}\right]\right\},$$
$$0 < \eta < \infty, \ -\frac{\pi}{2} < \phi_j < \frac{\pi}{2}, \quad j = 1, 2, \tag{2.1}$$

where

$$\chi(s) = \frac{(2\pi)^s}{\pi}\sin\left(\frac{\pi s}{2}\right)\Gamma(1-s), \tag{2.2}$$

with $\Gamma(s)$, $s \in \mathbb{C}$, *denoting the Gamma function, and where the contours of the first and second integrals in the rhs of (2.1) are the rays from* $\eta\exp(-i\pi/2)$ *to* $\infty\exp(i\phi_1)$ *and from* $\eta\exp(i\pi/2)$ *to* $\infty\exp(i\phi_2)$ *respectively. An equivalent formula is*

$$\zeta(1-s) = \sum_{n=1}^{\left[\frac{\eta}{2\pi}\right]} n^{s-1} + \frac{1}{(2\pi)^s}\left[-\frac{\eta^s}{s} + \left(e^{\frac{i\pi s}{2}}\int_{\eta e^{-\frac{i\pi}{2}}}^{\infty e^{i\phi_1}} + e^{-\frac{i\pi s}{2}}\int_{\eta e^{\frac{i\pi}{2}}}^{\infty e^{i\phi_2}}\right)\frac{e^{-z}z^{s-1}dz}{1-e^{-z}}\right],$$
$$0 < \eta < \infty, \ -\frac{\pi}{2} < \phi_j < \frac{\pi}{2}, \quad j = 1, 2. \tag{2.3}$$

Proof. The Riemann function $\zeta(s)$ can be defined by

$$\zeta(s) = \frac{\Gamma(1-s)}{2i\pi}\int_{H_\alpha}\frac{z^{s-1}}{e^{-z}-1}dz, \qquad s \in \mathbb{C}, \tag{2.4}$$

where H_α denotes the so-called Hankel contour, which is the union of the following curves in the complex z-plane cut along the negative real axis (see figure 1):

$$H_\alpha = \left\{\alpha e^{i\theta}, -\pi < \theta < \pi\right\} \cup \left\{|z|e^{-i\pi}, \quad \alpha < |z| < \infty\right\} \cup \left\{|z|e^{i\pi}, \quad \alpha < |z| < \infty\right\},$$
$$0 < \alpha < 2\pi. \tag{2.5}$$

We will first derive equation (2.1) in the particular case $0 < \eta < 2\pi$. In this case replacing η with α we find

$$\zeta(s) = \frac{\chi(s)}{(2\pi)^s}\left\{-\frac{\alpha^s}{s} + \left(e^{\frac{i\pi s}{2}}\int_{\alpha e^{-\frac{i\pi}{2}}}^{\infty e^{i\phi_1}} + e^{-\frac{i\pi s}{2}}\int_{\alpha e^{\frac{i\pi}{2}}}^{\infty e^{i\phi_2}}\right)\frac{e^{-z}z^{s-1}}{1-e^{-z}}dz\right\},$$
$$0 < \alpha < 2\pi; \quad \frac{-\pi}{2} < \phi_j < \frac{\pi}{2}, \quad j = 1, 2. \tag{2.6}$$



Figure 1 *The decomposition of H_α into $L_1 + L_2 + L_3$.*

We decompose the contour H_α into the union of three different contours (see figure 1), namely

$$H_\alpha = L_1 \cup L_2 \cup L_3,$$

where

$$\begin{aligned} L_1 &= \left\{\alpha e^{i\theta}, \ \ \frac{\pi}{2} < \theta < \pi\right\} \cup \{|z|e^{i\pi}, \ \ \alpha < |z| < \infty)\}, \\ L_2 &= \{|z|e^{-i\pi}, \ \ \alpha < |z| < \infty\} \cup \left\{\alpha e^{i\theta}, \ \ -\pi < \theta < -\frac{\pi}{2}\right\}, \\ L_3 &= \left\{\alpha e^{i\theta}, \ \ -\frac{\pi}{2} < \theta < \frac{\pi}{2}\right\}. \end{aligned} \tag{2.7}$$

The integral along L_1 can be written as follows:

$$\int_{L_1} \frac{z^{s-1}}{e^{-z}-1} dz = \int_{\alpha e^{\frac{i\pi}{2}}}^{\infty e^{i\pi}} \frac{e^z}{1-e^z} z^{s-1} dz = e^{i\pi s} \int_{\alpha e^{-\frac{i\pi}{2}}}^{\infty} \frac{e^{-u} u^{s-1} du}{1-e^{-u}}, \tag{2.8}$$

where the curve L_1 is defined in (2.7) and the contours of the second and third integrals in (2.8) are the rays in the cut complex z-plane from $\alpha \exp(i\pi/2)$ to $\infty \exp(i\pi)$ and from $\alpha \exp(-i\pi/2)$ to ∞ respectively.

Indeed, in the domain enclosed by L_1 and by the ray from $\alpha \exp(i\pi/2)$ to $\infty \exp(i\pi)$, i.e. in the shaded domain 1 of figure 2, we have

$$z = re^{i\theta}, \quad \alpha \le r \le \infty, \quad \frac{\pi}{2} \le \theta \le \pi;$$

thus,

$$|e^z| = e^{r\cos\theta}, \quad \alpha \le r \le \infty, \quad 0 \le \cos\theta \le -1.$$



Figure 2 *The domains* 1*,* 2*, and* 3*.*

Hence, Cauchy's theorem implies the first equality of (2.8). Then, the substitution $u = z\exp(-i\pi)$, yields the second equality in (2.6). Similarly,

$$\int_{L_2} \frac{z^{s-1}}{e^{-z}-1}dz = \int_{\infty e^{-i\pi}}^{\alpha e^{-\frac{i\pi}{2}}} \frac{e^z}{1-e^z} z^{s-1}dz = e^{-i\pi s}\int_{\infty}^{\alpha e^{\frac{i\pi}{2}}} \frac{e^{-u}u^{s-1}}{1-e^{-u}}du, \tag{2.9}$$

where the curve L_2 is defined in (2.7) and the contours of the second and third integrals in (2.9) are the rays in the cut complex z-plane from $\infty\exp(-i\pi)$ to $\alpha\exp(-i\pi/2)$ and from ∞ to $\alpha\exp(i\pi/2)$ respectively.

Indeed, the first equality in (2.9) is the consequence of Cauchy's theorem applied in the domain enclosed by L_2 and by the ray from $\infty\exp(-i\pi)$ to $\alpha\exp(-i\pi/2)$ (i.e. in the shaded domain 2 of figure 2), whereas the second equality in (2.9) follows from the substitution $u = z\exp(i\pi)$.

The integral along L_3 can be written as follows:

$$\int_{L_3} \frac{z^{s-1}}{e^{-z}-1}dz = -\left(\int_{\alpha e^{-\frac{i\pi}{2}}}^{\infty} + \int_{\infty}^{\alpha e^{\frac{i\pi}{2}}}\right) \frac{e^{-z}}{1-e^{-z}} z^{s-1}dz - 2i\sin\left(\frac{\pi s}{2}\right)\frac{\alpha^s}{s}, \tag{2.10}$$

where the curve L_3 is defined in (2.7) and the contours in the first and second integrals in (2.10) are the rays in the cut complex z-plane from $\alpha\exp(-i\pi/2)$ to ∞ and from ∞ to $\alpha\exp(i\pi/2)$.

Indeed,

$$\int_{L_3} \frac{z^{s-1}}{e^{-z}-1}dz = -\int_{L_3} \frac{e^{-z}}{1-e^{-z}} z^{s-1}dz - \int_{L_3} z^{s-1}dz. \tag{2.11}$$

The second term in the rhs of (2.11) yields the second term in the rhs of (2.10). Furthermore, in the domain enclosed by L_3 and by the two rays $(\alpha\exp(-i\pi/2),\infty)$ and $(\infty,\alpha\exp(i\pi/2))$, i.e. in the shaded domain 3 of figure 2, we have

$$z = re^{i\theta}, \quad \alpha \le r \le \infty, \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2};$$

thus

$$|e^{-z}| = e^{-r\cos\theta}, \quad \alpha \le r \le \infty, \quad \cos\theta \ge 0.$$

Hence, Cauchy's theorem implies that the first term of the rhs of (2.11) equals the first term in the rhs of (2.10).



Figure 3 *The curves* C_α^η, $C_{-\eta}^{-\alpha}$ *and* $\hat{C}_\eta^\alpha$*.*

Adding equations (2.8)-(2.10) we find equation (2.6) but with $\phi_1 = \phi_2 = 0$. However, Cauchy's theorem implies that the rays $(\alpha \exp(-i\pi/2), \infty)$ and $(\infty, \alpha \exp(i\pi/2))$ can be replaced with the rays $(\alpha \exp(-i\pi/2), \infty \exp(i\phi_1))$ and $(\infty \exp(i\phi_2), \alpha \exp(i\pi/2))$, respectively, and then we find (2.6).

In order to derive equation (2.1) with $2\pi < \eta < \infty$, we introduce the curves C_α^η, $C_{-\eta}^{-\alpha}$ and $\hat{C}_\eta^\alpha$, which are the following semi-circles in the complex z-plane, see figure 3:

$$C_\alpha^\eta = \left\{ \frac{i(\eta+\alpha)}{2} + \frac{(\eta-\alpha)}{2} e^{i\theta}, \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2} \right\}, \tag{2.12a}$$

$$C_{-\eta}^{-\alpha} = \left\{ \frac{-i(\eta+\alpha)}{2} + \frac{(\eta-\alpha)}{2} e^{i\theta}, \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2} \right\}, \tag{2.12b}$$

$$\hat{C}_\eta^\alpha = \left\{ \frac{i(\eta+\alpha)}{2} + \frac{(\eta-\alpha)}{2} e^{i\theta}, \quad \left(-\pi < \theta < \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2} < \theta < \pi\right) \right\}. \tag{2.12c}$$

The second integral in the rhs of equation (2.6) can be rewritten as an integral along the curve C_α^η plus an integral along the ray $(\eta \exp(i\pi/2), \infty \exp(i\phi_1))$. Similarly, the third integral in the rhs of equation (2.6) can be rewritten as an integral along the curve $-C_{-\eta}^{-\alpha}$ plus an integral along the ray $(\eta \exp(i\pi/2), \infty \exp(i\phi_2))$. Hence, the sum of the second and third integrals in the rhs of equation (2.6) yield the sum of the second and third integral in equation (2.1) plus the sum of the following two integrals:

$$I_1 = -e^{\frac{i\pi s}{2}} \int_{C_{-\eta}^{-\alpha}} \frac{e^{-z}}{1-e^{-z}} z^{s-1} dz, \quad I_2 = e^{-\frac{i\pi s}{2}} \int_{C_\alpha^\eta} \frac{e^{-z}}{1-e^{-z}} z^{s-1} dz. \tag{2.13}$$

Making the change of variables $u = z \exp(i\pi)$ in the integral I_1, we find

$$I_1 = e^{-\frac{i\pi s}{2}} \int_{\hat{C}_\eta^\alpha} \frac{1}{1-e^{-u}} u^{s-1} du = e^{-\frac{i\pi s}{2}} \int_{\hat{C}_\eta^\alpha} \frac{e^{-u}}{1-e^{-u}} u^{s-1} du + e^{-\frac{i\pi s}{2}} \int_{\hat{C}_\eta^\alpha} u^{s-1} du$$

$$= e^{-\frac{i\pi s}{2}} \int_{\hat{C}_\eta^\alpha} \frac{e^{-u}}{1-e^{-u}} u^{s-1} du + \frac{1}{s}(\alpha^s - \eta^s). \tag{2.14}$$

The sum of I_2 and of the first term in the rhs of equation (2.14) can be evaluated using Cauchy's theorem:

$$I_1 + I_2 = (2\pi)^s \sum_{m=1}^{\left[\frac{\eta}{2\pi}\right]} m^{s-1} + \frac{1}{s}(\alpha^s - \eta^s).$$

Hence equation (2.1) follows.

Multiplying equation (2.1) by $\chi(1-s)$ and using the identity

$$\chi(s)\chi(1-s) = 1, \quad s \in \mathbb{C}, \tag{2.15}$$

together with the functional equation

$$\zeta(1-s) = \chi(1-s)\zeta(s), \quad s \in \mathbb{C}, \tag{2.16}$$

we find the equation (2.3). **QED**

3 The Asymptotics of the Riemann Zeta Function for $t \leq \eta < \infty$

We are mainly interested in the behavior of $\zeta(s)$ in the critical strip $0 \leq \operatorname{Re} s \leq 1$, thus we will henceforth assume that $\sigma \in [0,1]$. It is possible to extend the results of this section to the case when σ belongs to any bounded interval, but the proofs are more complicated (e.g. equation (3.12) involves powers of σ that cannot be neglected if $\sigma > 1$).

Theorem 3.1 (The asymptotics to all orders for the case $(1+\epsilon)t < \eta$) *Let $\zeta(s)$, $s = \sigma + it$, $\sigma, t \in \mathbb{R}$, denote the Riemann zeta function. Then,*

$$\begin{aligned}\zeta(s) = {} & \sum_{n=1}^{\left[\frac{\eta}{2\pi}\right]} n^{-s} - \frac{1}{1-s}\left(\frac{\eta}{2\pi}\right)^{1-s} \\ & + \frac{e^{-\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}} \sum_{n=1}^{\infty}\sum_{j=0}^{N-1} e^{-nz-it\ln z}\left(\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\right)^j \frac{z^{-\sigma}}{n+\frac{it}{z}}\Bigg|_{z=i\eta} \\ & + \frac{e^{\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}} \sum_{n=1}^{\infty}\sum_{j=0}^{N-1} e^{-nz-it\ln z}\left(\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\right)^j \frac{z^{-\sigma}}{n+\frac{it}{z}}\Bigg|_{z=-i\eta} \\ & + O\left((2N+1)!!N\left(\frac{1+\epsilon}{\epsilon}\right)^{2(N+1)}\eta^{-\sigma-N}\right),\end{aligned} \tag{3.1}$$

$$(1+\epsilon)t < \eta < \infty, \quad \epsilon > 0, \quad 0 \le \sigma \le 1, \quad N \ge 2, \quad t \to \infty,$$

where the error term is uniform for all $\eta, \epsilon, \sigma, N$ *in the above ranges and* $(2N+1)!!$ *is defined by*

$$(2N+1)!! = 1 \cdot 3 \cdot 5 \cdot \dots \cdot (2N-1)(2N+1).$$

For $N = 3$ *equation (3.1) simplifies to*

$$\begin{aligned}\zeta(s) = {} & \sum_{n=1}^{[\frac{\eta}{2\pi}]} n^{-s} - \frac{1}{1-s}\Big(\frac{\eta}{2\pi}\Big)^{1-s} \\ & + \frac{2i\eta^{-s}}{(2\pi)^{1-s}}\bigg\{-i\arg(1-e^{i\eta}) + \frac{t-i\sigma}{\eta}\operatorname{Re}\operatorname{Li}_2(e^{i\eta}) \\ & \qquad + \frac{1}{\eta^2}\big[it^2 - 3i\sigma t - (\sigma-1)t - i\sigma(\sigma+1)\big]\operatorname{Im}\operatorname{Li}_3(e^{i\eta})\bigg\} \\ & + O\bigg(\frac{1}{\eta^{3+\sigma}}\Big(t^3 + \Big(\frac{1+\epsilon}{\epsilon}\Big)^8\Big)\bigg), \\ & \qquad (1+\epsilon)t < \eta < \infty, \quad \epsilon > 0, \quad 0 \le \sigma \le 1, \quad t \to \infty,\end{aligned} \tag{3.2}$$

where the error term is uniform for all η, ϵ, σ *in the above ranges and the polylogarithm* $\operatorname{Li}_m(z)$ *is defined by*

$$\operatorname{Li}_m(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^m}, \qquad m \ge 1. \tag{3.3}$$

Similarly, it is straightforward for any $N \ge 4$ *to derive an asymptotic formula for* $\zeta(s)$ *analogous to (3.2) with an error term of order*

$$O\bigg(\frac{1}{\eta^{N+\sigma}}\Big(t^N + \Big(\frac{1+\epsilon}{\epsilon}\Big)^{2(N+1)}\Big)\bigg).$$

Proof. Suppose first that $0 < t \le \eta < \infty$, $\epsilon > 0$, $0 \le \sigma \le 1$, $N \ge 2$. All error terms of the form $O(\cdot)$ will be uniform with respect to $\eta, \epsilon, \sigma, N$ unless otherwise specified. The proof is based on equation (2.3), i.e.

$$\zeta(1-s) = \sum_{n=1}^{[\frac{\eta}{2\pi}]} n^{s-1} - \frac{1}{s}\Big(\frac{\eta}{2\pi}\Big)^{s} + G_L(t,\sigma;\eta) + G_U(t,\sigma;\eta), \qquad 0 < \eta < \infty, \tag{3.4}$$

where G_L and G_U are defined by

$$G_L(t,\sigma;\eta) = \frac{e^{\frac{i\pi s}{2}}}{(2\pi)^s}\int_{-i\eta}^{\infty e^{i\phi_1}} \frac{e^{-z}z^{s-1}dz}{1-e^{-z}}, \qquad -\frac{\pi}{2} < \phi_1 < \frac{\pi}{2}, \quad 0 < \eta < \infty, \tag{3.5}$$

and

$$G_U(t,\sigma;\eta) = \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s}\int_{i\eta}^{\infty e^{i\phi_2}} \frac{e^{-z}z^{s-1}dz}{1-e^{-z}}, \qquad -\frac{\pi}{2} < \phi_2 < \frac{\pi}{2}, \quad 0 < \eta < \infty. \tag{3.6}$$

The asymptotics of G_L

Using the expansion

$$\frac{1}{1-e^{-z}} = \sum_{n=0}^{\infty} e^{-nz}, \qquad \operatorname{Re} z > 0, \tag{3.7}$$

in the definition (3.5) of G_L, we find

$$G_L(t,\sigma;\eta) = \frac{e^{\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} \int_{-i\eta}^{\infty e^{i\phi_1}} e^{-nz} z^{s-1} dz, \qquad -\frac{\pi}{2} < \phi_1 < \frac{\pi}{2}, \quad 0 < \eta < \infty. \tag{3.8}$$

The interchange of the integration and the summation in (3.8) is allowed because the sum on the rhs of (3.7) converges absolutely and uniformly in $\operatorname{Re} z \geq \delta$ for any $\delta > 0$, and because by assumption $\eta \notin 2\pi\mathbb{Z}$. Integration by parts shows that the integral in (3.8) can be written as

$$\begin{aligned}\int_{-i\eta}^{\infty e^{i\phi_1}} e^{-nz} z^{s-1} dz = {} & \sum_{j=0}^{N-1} e^{-nz+it\ln z} \left(\frac{1}{n-\frac{it}{z}} \frac{d}{dz} \right)^j \frac{z^{\sigma-1}}{n-\frac{it}{z}} \Bigg|_{z=-i\eta} \\ & + \int_{-i\eta}^{\infty e^{i\phi_1}} e^{-nz+it\ln z} D_N dz,\end{aligned} \tag{3.9}$$

where $D_N = D_N(z,n,\sigma,t)$ is short-hand notation for

$$D_N := \left(\frac{d}{dz} \frac{1}{n-\frac{it}{z}} \right)^N z^{\sigma-1}. \tag{3.10}$$

We claim that

$$\frac{e^{\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} \int_{-i\eta}^{\infty e^{i\phi_1}} e^{-nz+it\ln z} D_N dz = O\Big((2N-1)!!(N+1)^2 \eta^{\sigma-1-N} \Big). \tag{3.11}$$

In order to derive equation (3.11) we first note that we can write D_N in the form

$$D_N = \sum_{b=0}^{N} \sum_{c=0}^{N} A_{bc}^{(N)} z^{\sigma-1} \frac{t^b (nz)^{N-b} \sigma^c}{(nz-it)^{2N}} \tag{3.12}$$

where $A_{bc}^{(N)}$, $b = 0, \dots, N$, $c = 0, \dots, N$, are integers which satisfy

$$|A_{bc}^{(N)}| \leq (2N-1)!! = 1 \cdot 3 \cdot 5 \cdot \dots \cdot (2N-1), \qquad b = 0, \dots, N, \quad c = 0, \dots, N.$$

Indeed, if D_N can be written as in (3.12), then the computation

$$\frac{d}{dz}\left(\frac{1}{n-\frac{it}{z}} A_{bc}^{(N)} z^{\sigma-1} \frac{t^b (nz)^{N-b} \sigma^c}{(nz-it)^{2N}} \right)$$

$$= A_{bc}^{(N)} z^{\sigma-1} \frac{t^b (nz)^{N-b} \sigma^c}{(nz-it)^{2(N+1)}} [it(b-N) - it\sigma - nz(1+b+N) + nz\sigma)]$$

shows that D_{N+1} also can be written as in (3.12).

In order to estimate the lhs of (3.11), we choose $\phi_1 = 0$. Then, since $t \leq |z|$ and $|n - \frac{it}{z}| \geq n$ for all z on the contour, we can estimate

$$\begin{aligned} D_N &= O\biggl((2N-1)!! |z|^{\sigma-1} \sum_{b=0}^{N} \sum_{c=0}^{N} \biggl| \left(\frac{t}{nz}\right)^b \frac{(nz)^N \sigma^c}{z^{2N}(n - \frac{it}{z})^{2N}} \biggr| \biggr) \\ &= O\biggl((2N-1)!! |z|^{\sigma-1-N} (N+1)^2 \frac{1}{n^N} \biggr), \qquad z = u - i\eta, \quad u \in [0, \infty). \end{aligned}$$

Thus, the lhs of (3.11) is

$$\begin{aligned} &O\biggl(e^{-\frac{\pi t}{2}} \sum_{n=1}^{\infty} \int_{-i\eta}^{\infty} |e^{-nz} e^{it\log z}| (2N-1)!! (N+1)^2 |z|^{\sigma-1-N} \frac{1}{n^N} dz \biggr) \\ &= O\biggl((2N-1)!! (N+1)^2 \sum_{n=1}^{\infty} \frac{1}{n^N} \int_{-i\eta}^{\infty} e^{-n \operatorname{Re} z} |z|^{\sigma-1-N} dz \biggr) \\ &= O\biggl((2N-1)!! (N+1)^2 \eta^{\sigma-1-N} \sum_{n=1}^{\infty} \frac{1}{n^N} \int_{0}^{\infty} e^{-nu} du \biggr) \\ &= O\biggl((2N-1)!! (N+1)^2 \eta^{\sigma-1-N} \sum_{n=1}^{\infty} \frac{1}{n^{N+1}} \biggr) \\ &= O\biggl((2N-1)!! (N+1)^2 \eta^{\sigma-1-N} \biggr). \end{aligned}$$

This proves (3.11).

Equations (3.8), (3.9), and (3.11) imply that G_L satisfies

$$\begin{aligned} G_L(t,\sigma;\eta) = \frac{e^{\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} \sum_{j=0}^{N-1} e^{-nz+it\ln z} \left(\frac{1}{n - \frac{it}{z}} \frac{d}{dz} \right)^j \frac{z^{\sigma-1}}{n - \frac{it}{z}} \bigg|_{z=-i\eta} \\ + O\biggl((2N-1)!! (N+1)^2 \eta^{\sigma-1-N} \biggr), \qquad t \leq \eta < \infty, \quad t \to \infty. \end{aligned} \tag{3.13}$$

The asymptotics of G_U

We now let $\epsilon > 0$ and suppose that $0 < (1+\epsilon)t \leq \eta < \infty$. In analogy with equations (3.8) and (3.9) we now find

$$G_U(t,\sigma;\eta) = \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} \int_{i\eta}^{\infty e^{i\phi_2}} e^{-nz} z^{s-1} dz, \qquad -\frac{\pi}{2} < \phi_2 < \frac{\pi}{2}, \quad 0 < \eta < \infty, \tag{3.14}$$

and

$$\int_{i\eta}^{\infty e^{i\phi_2}} e^{-nz}z^{s-1}dz = \sum_{j=0}^{N-1} e^{-nz+it\ln z}\left(\frac{1}{n-\frac{it}{z}}\frac{d}{dz}\right)^j \frac{z^{\sigma-1}}{n-\frac{it}{z}}\bigg|_{z=i\eta} + \int_{i\eta}^{\infty e^{i\phi_2}} e^{-nz+it\ln z}D_N dz, \tag{3.15}$$

where D_N is defined by (3.10). We claim that

$$\frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s}\sum_{n=1}^{\infty}\int_{i\eta}^{\infty e^{i\phi_2}} e^{-nz+it\ln z}D_N dz = O\bigg((2N-1)!!N\Big(\frac{1+\epsilon}{\epsilon}\Big)^{2N}\eta^{\sigma-N}\bigg). \tag{3.16}$$

Indeed, since D_N can be written in the form given by (3.12), Jordan's lemma implies that we may choose $\phi_2 = \pi/2$. Then, using the estimate

$$\begin{aligned}D_N &= O\bigg((2N-1)!!|z|^{\sigma-1}\sum_{b=0}^{N}\sum_{c=0}^{N}\bigg|\Big(\frac{t}{nz}\Big)^b\frac{(nz)^N\sigma^c}{z^{2N}(n-\frac{it}{z})^{2N}}\bigg|\bigg)\\ &= O\bigg((2N-1)!!|z|^{\sigma-1-N}(N+1)^2\frac{n^N}{(n-\frac{t}{\eta})^{2N}}\bigg), \qquad z=i\lambda, \quad \lambda\in[\eta,\infty),\end{aligned}$$

we find that the lhs of (3.16) is

$$\begin{aligned}&O\bigg(e^{\frac{\pi t}{2}}\sum_{n=1}^{\infty}\int_{i\eta}^{i\infty}|e^{-nz+it\log z}D_N|dz\bigg)\\ &\quad= O\bigg(e^{\frac{\pi t}{2}}\sum_{n=1}^{\infty}\int_{i\eta}^{i\infty}e^{-\frac{\pi t}{2}}(2N-1)!!|z|^{\sigma-1-N}(N+1)^2\frac{n^N}{(n-\frac{t}{\eta})^{2N}}dz\bigg)\\ &\quad= O\bigg((2N-1)!!(N+1)^2\int_{\eta}^{\infty}\lambda^{\sigma-1-N}d\lambda\sum_{n=1}^{\infty}\frac{n^N}{(n-\frac{1}{1+\epsilon})^{2N}}\bigg)\\ &\quad= O\bigg((2N-1)!!(N+1)^2\frac{\eta^{\sigma-N}}{\sigma-N}\Big(\frac{1+\epsilon}{\epsilon}\Big)^{2N}\bigg)\\ &\quad= O\bigg((2N-1)!!N\Big(\frac{1+\epsilon}{\epsilon}\Big)^{2N}\eta^{\sigma-N}\bigg).\end{aligned}$$

In the third equality above we have used the following estimates:

$$\begin{aligned}\sum_{n=1}^{\infty}\frac{n^N}{(n-\frac{1}{1+\epsilon})^{2N}} &= \frac{1}{(1-\frac{1}{1+\epsilon})^{2N}} + \sum_{n=2}^{\infty}e^{N(\ln n-2\ln(n-\frac{1}{1+\epsilon}))}\\ &= O\bigg(\Big(\frac{1+\epsilon}{\epsilon}\Big)^{2N} + \sum_{n=2}^{\infty}e^{N(\ln n-2\ln n+\frac{2}{1+\epsilon})}\bigg)\end{aligned} \tag{3.17}$$

$$= O\left(\left(\frac{1+\epsilon}{\epsilon}\right)^{2N} + e^{\frac{2N}{1+\epsilon}}\right) = O\left(\left(\frac{1+\epsilon}{\epsilon}\right)^{2N}\right), \qquad N \geq 2,$$

where the last step is a consequence of the fact that $e^{\frac{1}{1+\epsilon}} < \frac{1+\epsilon}{\epsilon}$ for $\epsilon > 0$. This proves (3.16).

We next claim that (3.16) can be modified as follows:

$$\frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} \int_{i\eta}^{\infty e^{i\phi_2}} e^{-nz+it\ln z} D_N dz \tag{3.18}$$
$$= O\left((2N+1)!!N\left(\frac{1+\epsilon}{\epsilon}\right)^{2(N+1)} \eta^{\sigma-N-1}\right).$$

Indeed, integration by parts shows that the lhs of (3.16) equals

$$\frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} e^{-nz+it\ln z} \frac{1}{n-\frac{it}{z}} D_N \bigg|_{z=i\eta} + \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} \int_{i\eta}^{\infty e^{i\phi_2}} e^{-nz+it\ln z} D_{N+1} dz.$$

By (3.16), the second term is

$$O\left((2N+1)!!(N+1)\left(\frac{1+\epsilon}{\epsilon}\right)^{2(N+1)} \eta^{\sigma-N-1}\right),$$

while, in view of (3.12) and (3.17), the first term satisfies

$$\frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} e^{-ni\eta+it\ln i\eta} \frac{1}{n-\frac{t}{\eta}} \sum_{b=0}^{N} \sum_{c=0}^{N} A_{bc}^{(N)} (i\eta)^{\sigma-1} \frac{t^b (ni\eta)^{N-b} \sigma^c}{(ni\eta - it)^{2N}}$$
$$= O\left(\eta^{\sigma-1-N} \sum_{b=0}^{N} \sum_{c=0}^{N} |A_{bc}^{(N)}| \sum_{n=1}^{\infty} \frac{t^b}{\eta^b} \frac{n^{N-b}}{(n-\frac{t}{\eta})^{2N+1}}\right)$$
$$= O\left(\eta^{\sigma-1-N} (2N-1)!!(N+1)^2 \sum_{n=1}^{\infty} \frac{n^N}{(n-\frac{1}{1+\epsilon})^{2N+1}}\right)$$
$$= O\left(\eta^{\sigma-1-N} (2N-1)!!(N+1)^2 \left(\frac{1+\epsilon}{\epsilon}\right)^{2N+1}\right).$$

This proves (3.18).

Equations (3.14), (3.15), and (3.18) imply that G_U satisfies

$$G_U(t,\sigma;\eta) = \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} \sum_{j=0}^{N-1} e^{-nz+it\ln z} \left(\frac{1}{n-\frac{it}{z}} \frac{d}{dz}\right)^j \frac{z^{\sigma-1}}{n-\frac{it}{z}} \bigg|_{z=i\eta} \tag{3.19}$$
$$+ O\left((2N+1)!!N\left(\frac{1+\epsilon}{\epsilon}\right)^{2(N+1)} \eta^{\sigma-N-1}\right), \qquad (1+\epsilon)t < \eta < \infty.$$

Proof of (3.1)

Substituting the expressions (3.13) and (3.19) for G_L and G_U into (3.4), we find

$$
\begin{aligned}
\zeta(1-s) = {} & \sum_{n=1}^{[\frac{\eta}{2\pi}]} n^{s-1} - \frac{1}{s}\left(\frac{\eta}{2\pi}\right)^{s} \\
& + \frac{e^{\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} \sum_{j=0}^{N-1} e^{-nz+it\ln z} \left(\frac{1}{n-\frac{it}{z}}\frac{d}{dz}\right)^{j} \frac{z^{\sigma-1}}{n-\frac{it}{z}}\Bigg|_{z=-i\eta} \\
& + \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} \sum_{j=0}^{N-1} e^{-nz+it\ln z} \left(\frac{1}{n-\frac{it}{z}}\frac{d}{dz}\right)^{j} \frac{z^{\sigma-1}}{n-\frac{it}{z}}\Bigg|_{z=i\eta} \\
& + O\Bigg((2N+1)!!N\Big(\frac{1+\epsilon}{\epsilon}\Big)^{2(N+1)}\eta^{\sigma-N-1}\Bigg), \\
& \qquad\qquad\qquad\qquad (1+\epsilon)t < \eta < \infty, \quad t \to \infty.
\end{aligned}
$$

Replacing σ by $1-\sigma$ and taking the complex conjugate of the resulting equation, we find (3.1).

Proof of (3.2)

Letting $N=3$ in (3.1), we find

$$
\begin{aligned}
\zeta(s) = {} & \sum_{n=1}^{[\frac{\eta}{2\pi}]} n^{-s} - \frac{1}{1-s}\left(\frac{\eta}{2\pi}\right)^{1-s} \\
& + \frac{e^{-\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}}(i\eta)^{-s} \sum_{n=1}^{\infty} e^{-i\eta n} \Bigg\{ \frac{1}{n}\frac{1}{1+q} + \frac{1}{\eta n^2}\frac{i(q(\sigma-1)+\sigma)}{(1+q)^3} \\
& \qquad + \frac{1}{\eta^2 n^3}\frac{-q^2(\sigma-1)^2 + q\left(-2\sigma^2+\sigma+2\right) - \sigma(\sigma+1)}{(1+q)^5} \Bigg\} \\
& + \frac{e^{\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}}(-i\eta)^{-s} \sum_{n=1}^{\infty} e^{i\eta n} \Bigg\{ \frac{1}{n}\frac{1}{1-q} + \frac{1}{\eta n^2}\frac{i(q(\sigma-1)-\sigma)}{(1-q)^3} \\
& \qquad - \frac{1}{\eta^2 n^3}\frac{q^2(\sigma-1)^2 + q\left(-2\sigma^2+\sigma+2\right) + \sigma(\sigma+1)}{(1-q)^5} \Bigg\} \\
& + O\Bigg(\eta^{-\sigma-3}\Big(\frac{1+\epsilon}{\epsilon}\Big)^{8}\Bigg), \qquad q = \frac{t}{n\eta}.
\end{aligned}
$$

The above sums can be expanded as series in q with coefficients expressible in terms of the polylogarithms $\mathrm{Li}_m(z)$ defined by (3.3). For example,

$$
\sum_{n=1}^{\infty} \frac{e^{i\eta n}}{n(1-q)} = \sum_{n=1}^{\infty} \frac{e^{i\eta n}}{n} \sum_{k=0}^{\infty} \Big(\frac{t}{n\eta}\Big)^{k} = \sum_{k=0}^{\infty} \left(\frac{t}{\eta}\right)^{k} \sum_{n=1}^{\infty} \frac{e^{i\eta n}}{n^{k+1}} = \sum_{k=0}^{\infty} \left(\frac{t}{\eta}\right)^{k} \mathrm{Li}_{k+1}\left(e^{i\eta}\right),
$$

where the interchange of the two sums can be justified by separating the terms where $k = 0$ and noticing that the remaining double sum is absolutely convergent. More precisely,

$$\begin{aligned}
\lim_{N\to\infty}\lim_{M\to\infty}\sum_{n=1}^{N}\frac{e^{i\eta n}}{n}\sum_{k=0}^{M}\left(\frac{t}{n\eta}\right)^{k} &= \lim_{N\to\infty}\left(\sum_{n=1}^{N}\frac{e^{i\eta n}}{n}+\lim_{M\to\infty}\sum_{n=1}^{N}\frac{e^{i\eta n}}{n}\sum_{k=1}^{M}\left(\frac{t}{n\eta}\right)^{k}\right)\\
&= \mathrm{Li}_1(e^{i\eta})+\sum_{n=1}^{\infty}\frac{e^{i\eta n}}{n}\sum_{k=1}^{\infty}\left(\frac{t}{n\eta}\right)^{k}\\
&= \mathrm{Li}_1(e^{i\eta})+\sum_{k=1}^{\infty}\sum_{n=1}^{\infty}\left(\frac{t}{\eta}\right)^{k}\frac{e^{i\eta n}}{n^{k+1}}\\
&= \sum_{k=0}^{\infty}\left(\frac{t}{\eta}\right)^{k}\mathrm{Li}_{k+1}\left(e^{i\eta}\right).
\end{aligned}$$

Similarly, for any $j \geq 0$ and $l, m \geq 1$,

$$\begin{aligned}
\sum_{n=1}^{\infty}\frac{e^{i\eta n}}{n^{l}}\frac{q^{j}}{(1-q)^{m}} &= \sum_{n=1}^{\infty}\frac{e^{i\eta n}}{n^{l}}\sum_{k=0}^{\infty}\binom{k+m-1}{k}\left(\frac{t}{n\eta}\right)^{k+j}\\
&= \sum_{k=0}^{\infty}\binom{k+m-1}{k}\left(\frac{t}{\eta}\right)^{k+j}\mathrm{Li}_{k+l+j}(e^{i\eta}).
\end{aligned}$$

Letting $\eta \to -\eta$ in this equation, we find

$$\sum_{n=1}^{\infty}\frac{e^{-i\eta n}}{n^{l}}\frac{(-q)^{j}}{(1+q)^{m}} = \sum_{k=0}^{\infty}\binom{k+m-1}{k}\left(-\frac{t}{\eta}\right)^{k+j}\mathrm{Li}_{k+l+j}(e^{-i\eta}).$$

This gives the following expression for the Riemann zeta function:

$$\begin{aligned}
\zeta(s) = \sum_{n=1}^{[\frac{\eta}{2\pi}]} n^{-s} &- \frac{1}{1-s}\left(\frac{\eta}{2\pi}\right)^{1-s}\\
&+ \frac{e^{-\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}}(i\eta)^{-s}\Bigg\{\sum_{k=0}^{\infty}\left(-\frac{t}{\eta}\right)^{k}\mathrm{Li}_{k+1}(e^{-i\eta})+\frac{i\sigma}{\eta}\sum_{k=0}^{\infty}\binom{k+2}{k}\left(-\frac{t}{\eta}\right)^{k}\mathrm{Li}_{k+2}(e^{-i\eta})\\
&\qquad - \frac{i(\sigma-1)}{\eta}\sum_{k=0}^{\infty}\binom{k+2}{k}\left(-\frac{t}{\eta}\right)^{k+1}\mathrm{Li}_{k+3}(e^{-i\eta})\\
&\qquad - \frac{\sigma(\sigma+1)}{\eta^{2}}\sum_{k=0}^{\infty}\binom{k+4}{k}\left(-\frac{t}{\eta}\right)^{k}\mathrm{Li}_{k+3}(e^{-i\eta})\\
&\qquad + \frac{2\sigma^{2}-\sigma-2}{\eta^{2}}\sum_{k=0}^{\infty}\binom{k+4}{k}\left(-\frac{t}{\eta}\right)^{k+1}\mathrm{Li}_{k+4}(e^{-i\eta})
\end{aligned}$$

$$
\begin{aligned}
&\qquad\qquad - \frac{(\sigma-1)^2}{\eta^2}\sum_{k=0}^{\infty}\binom{k+4}{k}\left(-\frac{t}{\eta}\right)^{k+2}\mathrm{Li}_{k+5}(e^{-i\eta})\Bigg\}\\
&+\frac{e^{\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}}(-i\eta)^{-s}\Bigg\{\sum_{k=0}^{\infty}\left(\frac{t}{\eta}\right)^{k}\mathrm{Li}_{k+1}(e^{i\eta})-\frac{i\sigma}{\eta}\sum_{k=0}^{\infty}\binom{k+2}{k}\left(\frac{t}{\eta}\right)^{k}\mathrm{Li}_{k+2}(e^{i\eta})\\
&\qquad\qquad+\frac{i(\sigma-1)}{\eta}\sum_{k=0}^{\infty}\binom{k+2}{k}\left(\frac{t}{\eta}\right)^{k+1}\mathrm{Li}_{k+3}(e^{i\eta})\\
&\qquad\qquad-\frac{\sigma(\sigma+1)}{\eta^2}\sum_{k=0}^{\infty}\binom{k+4}{k}\left(\frac{t}{\eta}\right)^{k}\mathrm{Li}_{k+3}(e^{i\eta})\\
&\qquad\qquad+\frac{2\sigma^2-\sigma-1}{\eta^2}\sum_{k=0}^{\infty}\binom{k+4}{k}\left(\frac{t}{\eta}\right)^{k+1}\mathrm{Li}_{k+4}(e^{i\eta})\\
&\qquad\qquad-\frac{(\sigma-1)^2}{\eta^2}\sum_{k=0}^{\infty}\binom{k+4}{k}\left(\frac{t}{\eta}\right)^{k+2}\mathrm{Li}_{k+5}(e^{i\eta})\Bigg\}\\
&+O\left(\eta^{-\sigma-3}\left(\frac{1+\epsilon}{\epsilon}\right)^{8}\right).
\end{aligned}
$$

By including only the first few terms in the sums over k, we find an expansion for $\zeta(s)$ in powers of $\frac{t}{\eta}$. For example, including only the terms of order larger than $O(\frac{t^3}{\eta^{3+\sigma}})$ we have

$$
\begin{aligned}
\zeta(s)={}&\sum_{n=1}^{[\frac{\eta}{2\pi}]}n^{-s}-\frac{1}{1-s}\left(\frac{\eta}{2\pi}\right)^{1-s}\\
&-\frac{i\eta^{-s}}{(2\pi)^{1-s}}\Bigg\{\mathrm{Li}_1(e^{-i\eta})-\frac{t}{\eta}\mathrm{Li}_2(e^{-i\eta})+\left(\frac{t}{\eta}\right)^2\mathrm{Li}_3(e^{-i\eta})\\
&+\frac{i\sigma}{\eta}\mathrm{Li}_2(e^{-i\eta})-\frac{3\sigma t}{\eta^2}\mathrm{Li}_3(e^{-i\eta})\\
&+\frac{i(\sigma-1)t}{\eta^2}\mathrm{Li}_3(e^{-i\eta})-\frac{\sigma(\sigma+1)}{\eta^2}\mathrm{Li}_3(e^{-i\eta})\Bigg\}\\
&+\frac{i\eta^{-s}}{(2\pi)^{1-s}}\Bigg\{\mathrm{Li}_1(e^{i\eta})+\frac{t}{\eta}\mathrm{Li}_2(e^{i\eta})+\left(\frac{t}{\eta}\right)^2\mathrm{Li}_3(e^{i\eta})\\
&-\frac{i\sigma}{\eta}\mathrm{Li}_2(e^{i\eta})-\frac{3\sigma t}{\eta^2}\mathrm{Li}_3(e^{i\eta})\\
&+\frac{i(\sigma-1)t}{\eta^2}\mathrm{Li}_3(e^{i\eta})-\frac{\sigma(\sigma+1)}{\eta^2}\mathrm{Li}_3(e^{i\eta})\Bigg\}\\
&+O\left(\frac{1}{\eta^{3+\sigma}}\left(t^3+\left(\frac{1+\epsilon}{\epsilon}\right)^{8}\right)\right).
\end{aligned}
$$

After simplification we find

$$\begin{aligned}\zeta(s) = & \sum_{n=1}^{[\frac{\eta}{2\pi}]} n^{-s} - \frac{1}{1-s}\left(\frac{\eta}{2\pi}\right)^{1-s} \\ & + \frac{2i\eta^{-s}}{(2\pi)^{1-s}}\Bigg\{ i\mathrm{Im}\,\mathrm{Li}_1(e^{i\eta}) + \frac{t}{\eta}\mathrm{Re}\,\mathrm{Li}_2(e^{i\eta}) + i\left(\frac{t}{\eta}\right)^2 \mathrm{Im}\,\mathrm{Li}_3(e^{i\eta}) \\ & \qquad - \frac{i\sigma}{\eta}\mathrm{Re}\,\mathrm{Li}_2(e^{i\eta}) - \frac{3i\sigma t}{\eta^2}\mathrm{Im}\,\mathrm{Li}_3(e^{i\eta}) \\ & \qquad - \frac{(\sigma-1)t}{\eta^2}\mathrm{Im}\,\mathrm{Li}_3(e^{i\eta}) - \frac{i\sigma(\sigma+1)}{\eta^2}\mathrm{Im}\,\mathrm{Li}_3(e^{i\eta})\Bigg\} \\ & + O\left(\frac{1}{\eta^{3+\sigma}}\left(t^3 + \left(\frac{1+\epsilon}{\epsilon}\right)^8\right)\right),\end{aligned}$$

The identity $\mathrm{Li}_1\left(e^{i\eta}\right) = -\log(1-e^{i\eta})$ now yields (3.2). **QED**

Theorem 3.2 (The asymptotic expansion to all orders for the case $t = \eta$) *Let $\zeta(s)$, $s = \sigma + it$, $\sigma, t \in \mathbb{R}$, denote the Riemann zeta function. Then,*

$$\begin{aligned}\zeta(s) = & \sum_{n=1}^{[\frac{t}{2\pi}]} n^{-s} - \frac{1}{1-s}\left(\frac{t}{2\pi}\right)^{1-s} \\ & + \frac{e^{-\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}} \sum_{n=1}^{\infty}\sum_{j=0}^{N-1} e^{-nz-it\ln z}\left(\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\right)^j \frac{z^{-\sigma}}{n+\frac{it}{z}}\Bigg|_{z=it} \\ & + \frac{e^{\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}} \sum_{n=2}^{\infty}\sum_{j=0}^{N-1} e^{-nz-it\ln z}\left(\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\right)^j \frac{z^{-\sigma}}{n+\frac{it}{z}}\Bigg|_{z=-it} \\ & + \left(\frac{t}{2\pi}\right)^{1-s} e^{it} \sum_{k=0}^{2N} \frac{\overline{c_k(1-\sigma)}\Gamma(\frac{k+1}{2})}{t^{\frac{k+1}{2}}} + O\left((2N+1)!!Ne^{2(N+1)}t^{-\sigma-N}\right), \\ & \qquad 0 \le \sigma \le 1, \quad N \ge 2, \quad t \to \infty,\end{aligned} \tag{3.20}$$

where the error term is uniform for all σ, N in the above ranges and the coefficients $c_k(\sigma)$ are given by equation (3.30) below. The first few of the c_k's are given by

$$\begin{aligned} c_0(\sigma) &= \frac{1-i}{2}, \\ c_1(\sigma) &= \frac{i}{3} - i\sigma, \\ c_2(\sigma) &= -\frac{1+i}{12}\left(6\sigma^2 - 6\sigma + 1\right), \\ c_3(\sigma) &= \frac{1}{135}\left(-45\sigma^3 + 90\sigma^2 - 45\sigma + 4\right), \end{aligned} \tag{3.21}$$

$$
\begin{aligned}
c_4(\sigma) &= \frac{i-1}{432}\left(36\sigma^4 - 120\sigma^3 + 120\sigma^2 - 36\sigma + 1\right),\\
c_5(\sigma) &= \frac{i}{5670}\left(189\sigma^5 - 945\sigma^4 + 1575\sigma^3 - 987\sigma^2 + 168\sigma + 8\right),\\
c_6(\sigma) &= \frac{1+i}{194400}\left(1080\sigma^6 - 7560\sigma^5 + 18900\sigma^4 - 20160\sigma^3 + 8190\sigma^2 - 450\sigma - 139\right).
\end{aligned}
$$

For $N = 3$ equation (3.20) simplifies to

$$
\begin{aligned}
\zeta(s) = &\sum_{n=1}^{[\frac{t}{2\pi}]} n^{-s} - \frac{1}{1-s}\left(\frac{t}{2\pi}\right)^{1-s} \qquad (3.22)\\
&- \frac{2it^{-s}e^{it}}{(2\pi)^{1-s}}\Bigg\{\frac{e^{-it}}{2t^2}\left[-t^2 - it(\sigma-1) + (\sigma-1)^2\right] - i\mathrm{Im}\,\mathrm{Li}_1(e^{it}) + \frac{i\sigma}{t}\mathrm{Re}\,\mathrm{Li}_2(e^{it})\\
&\qquad + \frac{it+\sigma+\sigma^2}{t^2} i\mathrm{Im}\,\mathrm{Li}_3(e^{it}) + \frac{2+3\sigma}{t^2}\mathrm{Re}\,\mathrm{Li}_4(e^{it}) + \frac{3i}{t^2}\mathrm{Im}\,\mathrm{Li}_5(e^{it})\Bigg\}\\
&+ \frac{t^{-s}e^{it}}{(2\pi)^{1-s}}\Bigg\{\frac{1+i}{2}\sqrt{\pi t} - \frac{1}{3}i(3\sigma-2)\\
&\qquad + \frac{i-1}{24\sqrt{t}}\sqrt{\pi}\left(6\sigma^2 - 6\sigma + 1\right) + \frac{1}{135t}\left(45\sigma^3 - 45\sigma^2 + 4\right)\\
&\qquad - \frac{1+i}{576t^{\frac{3}{2}}}\sqrt{\pi}\left(36\sigma^4 - 24\sigma^3 - 24\sigma^2 + 12\sigma + 1\right)\\
&\qquad + \frac{i}{2835t^2}\left(189\sigma^5 - 315\sigma^3 + 42\sigma^2 + 84\sigma - 8\right)\\
&\qquad + \frac{1-i}{103680t^{\frac{5}{2}}}\sqrt{\pi}\big(1080\sigma^6 + 1080\sigma^5 - 2700\sigma^4 - 1440\sigma^3\\
&\qquad + 1710\sigma^2 + 270\sigma - 139\big)\Bigg\} + O\big(t^{-\sigma-3}\big), \qquad 0 \le \sigma \le 1, \quad t \to \infty,
\end{aligned}
$$

where the error term is uniform for all σ in the above range and the polylogarithm $\mathrm{Li}_m(z)$, $m \ge 1$, is defined by (3.3). Similarly, it is straightforward for any $N \ge 4$ to derive an asymptotic formula for $\zeta(s)$ analogous to (3.22) with an error term of order $O\big(t^{-\sigma-N}\big)$.

Proof. Setting $\eta = t$ in (3.4), we find

$$
\zeta(1-s) = \sum_{n=1}^{[\frac{t}{2\pi}]} n^{s-1} - \frac{1}{s}\left(\frac{t}{2\pi}\right)^{s} + G_L(t,\sigma;t) + G_U(t,\sigma;t), \qquad 0 < t < \infty. \qquad (3.23)
$$

Equation (3.13) is valid also when $\eta = t$ and gives the asymptotics of $G_L(t,\sigma;t)$. On the other hand, using the expansion (3.7), we write

$$
G_U(t,\sigma;t) = G_U^{(1)}(t,\sigma;t) + G_U^{(2)}(t,\sigma;t),
$$

where

$$G_U^{(1)}(t,\sigma;t) = \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s}\int_{it}^{\infty e^{i\phi_2}} e^{-z}z^{s-1}dz, \qquad -\frac{\pi}{2} < \phi_2 < \frac{\pi}{2}, \quad 0 < t < \infty \tag{3.24}$$

and

$$G_U^{(2)}(t,\sigma;t) = \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s}\sum_{n=2}^{\infty}\int_{it}^{\infty e^{i\phi_2}} e^{-nz}z^{s-1}dz, \qquad -\frac{\pi}{2} < \phi_2 < \frac{\pi}{2}, \quad 0 < t < \infty. \tag{3.25}$$

The asymptotics of $G_U^{(2)}$ can be found using integration by parts, whereas the asymptotics of $G_U^{(1)}$ will be computed by considering the critical point at $z = it$.

The asymptotics of $G_U^{(2)}$

Repeating the steps that led to (3.19) but with the sum over n only going from 2 to ∞ and with $t = \eta$, we find the following analog of (3.19):

$$\begin{aligned} G_U^{(2)}(t,\sigma;t) = \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s}\sum_{n=2}^{\infty}\sum_{j=0}^{N-1} e^{-nz+it\ln z}\left(\frac{1}{n-\frac{it}{z}}\frac{d}{dz}\right)^j \frac{z^{\sigma-1}}{n-\frac{it}{z}}\Bigg|_{z=it} \\ + O\bigg((2N+1)!!Ne^{2(N+1)}t^{\sigma-N-1}\bigg). \end{aligned} \tag{3.26}$$

The asymptotics of $G_U^{(1)}$

Letting in the definition (3.24) of $G_U^{(1)}$, $\phi_2 = 0$ and

$$z = it + t\rho, \qquad \rho \in [0,\infty),$$

we find

$$G_U^{(1)} = \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s}(it)^{s-1}te^{-it}\int_0^{\infty} e^{-t(\rho - i\log(1-i\rho))}(1-i\rho)^{\sigma-1}d\rho.$$

Letting

$$v = \rho - i\log(1-i\rho), \tag{3.27}$$

we find

$$G_U^{(1)} = \left(\frac{t}{2\pi}\right)^s e^{-it}\int_{\gamma} e^{-tv}\frac{(1-i\rho(v))^{\sigma}}{\rho(v)}dv,$$

where $\rho(v)$ is defined by inverting (3.27) and γ denotes the image of the contour $[0,\infty)$ under (3.27). In order to ascertain that the value of $\rho(v)$ is well-defined, we consider in detail the map ϕ defined by

$$\phi : \mathbb{C}\setminus[-i,-i\infty) \to \mathbb{C},$$

$$\rho \mapsto v = \rho - i \log(1 - i\rho),$$

where $[-i, -i\infty)$ is a branch cut and the principal branch is chosen for the logarithm, i.e.

$$v = \rho - i \ln|1 - i\rho| + \arg(1 - i\rho), \qquad \arg(1 - i\rho) \in (-\pi, \pi). \tag{3.28}$$

The function ϕ satisfies

$$\begin{aligned} &\phi(\rho) = 0 \quad \text{iff} \quad \rho = 0; \\ &\phi'(\rho) = \frac{\rho}{i + \rho} = 0 \quad \text{iff} \quad \rho = 0; \\ &v = \phi(\rho) = -\frac{i\rho^2}{2} + O(\rho^3), \qquad \rho \to 0. \end{aligned}$$

Moreover, we claim that ϕ maps the first quadrant of the complex ρ-plane bijectively onto the region delimited by the contour $\gamma = \phi([0, \infty))$ and the positive imaginary axis in the complex v-plane, see figure 4. Indeed, it is clear that ϕ maps $[0, i\infty)$ bijectively onto $[0, i\infty)$. Writing $\rho = \rho_1 + i\rho_2$, we have

$$\frac{\partial \mathrm{Re}\, \phi}{\partial \rho_1} = \frac{|\rho|^2 + \rho_2}{\rho_1^2 + (\rho_2 + 1)^2}, \qquad \frac{\partial \mathrm{Re}\, \phi}{\partial \rho_2} = \frac{\rho_1}{\rho_1^2 + (\rho_2 + 1)^2},$$

This shows that each level curve of the function $\mathrm{Re}\, \phi$ in the first quadrant intersects the positive ρ-axis in a unique point and that the vector

$$\big(-\rho_1, |\rho|^2 + \rho_2\big) \tag{3.29}$$

is tangent to the level curve passing through ρ. Since

$$\frac{\partial \mathrm{Im}\, \phi}{\partial \rho_1} = \frac{-\rho_1}{\rho_1^2 + (\rho_2 + 1)^2}, \qquad \frac{\partial \mathrm{Im}\, \phi}{\partial \rho_2} = \frac{|\rho|^2 + \rho_2}{\rho_1^2 + (\rho_2 + 1)^2},$$

we infer that the derivative of $\mathrm{Im}\, \phi$ in the direction of (3.29) equals $|\rho|^2$, showing that $\mathrm{Im}\, \phi$ is strictly increasing along each level curve. This proves the claim.

Thus, if v belongs to the region delimited by the contour $\gamma = \phi([0, \infty))$ and the positive imaginary axis, we may define $\rho(v)$ as the unique inverse image of v under ϕ which belongs to the first quadrant. Using analyticity to deform the contour γ to the positive real axis, we obtain

$$G_U^{(1)} = \left(\frac{t}{2\pi}\right)^s e^{-it} \int_0^\infty e^{-tv} \frac{(1 - i\rho(v))^\sigma}{\rho(v)} dv.$$

We claim that there exist coefficients $\{c_k(\sigma)\}_0^{N-1}$ such that

$$\frac{(1 - i\rho(v))^\sigma}{\rho(v)} = \frac{1}{\sqrt{v}} \bigg(\sum_{k=0}^{N-1} c_k(\sigma) v^{k/2} + r_N(\sqrt{v}) \bigg), \qquad v \geq 0, \tag{3.30}$$



Figure 4 *The function ϕ maps the indicated shaded areas in the complex ρ-plane bijectively onto the corresponding shaded areas in the v-plane.*

where $r_N(\sqrt{v})$ satisfies

$$\int_0^\infty e^{-tv}\frac{r_N(\sqrt{v})}{\sqrt{v}}dv = O\bigg(\frac{\Gamma(\frac{N+1}{2})}{t^{\frac{N+1}{2}}}\bigg). \tag{3.31}$$

Indeed, since ϕ has a double zero at $\rho = 0$, there exists a neighborhood V of $\rho = 0$ in the complex ρ-plane and a Riemann surface Σ which is a two-sheeted cover of V with local parameter $\lambda := \sqrt{v}$, such that the map $\rho \mapsto \sqrt{v}$ is bijective and holomorphic $V \to \Sigma$. The function ϕ is analytic away from $[-i, -i\infty)$ and $\phi'(\rho) \neq 0$ for all $\rho \neq 0$, thus the Riemann surface Σ and the map $\sqrt{v} \mapsto \rho$ can be analytically extended as long as $\phi^{-1}(v) \cap [-i, -i\infty) = \emptyset$. The function ϕ maps the segments $[-i+0, -i\infty+0)$ and $[-i-0, -i\infty-0)$ onto the lines $\operatorname{Re} v = -\pi$ and $\operatorname{Re} v = \pi$ respectively, hence $\sqrt{v} \mapsto \rho$ is well-defined and analytic for all $|v| < \pi$. Thus, the map

$$f : \lambda \mapsto \lambda\frac{(1-i\rho)^\sigma}{\rho} = \lambda\frac{e^{\sigma\log(1-i\rho)}}{\rho}$$

is holomorphic from the open disk $\{0 < |\lambda| < \sqrt{\pi}\} \subset \Sigma$ to $\mathbb{C}$. This implies that (3.30) holds with

$$c_k = \frac{f^{(k)}(0)}{k!}, \quad k = 0, \dots, N-1,$$

and with a function $r_N(\lambda)$ satisfying

$$r_N(\lambda) = \sum_{k=N}^\infty \frac{f^{(k)}(0)}{k!}\lambda^k, \qquad 0 \leq |\lambda| < \sqrt{\pi}.$$

In order to prove (3.31), it is sufficient to show that there exists a constant C independent of N such that

$$|r_N(\sqrt{v})| \leq Cv^{\frac{N}{2}}, \qquad v \geq 0, \quad N \geq 2. \tag{3.32}$$

Cauchy's estimates imply that the numbers $c_k = \frac{f^{(k)}(0)}{k!}$ are uniformly bounded for all $k \geq 0$. Thus, equations (3.28) and (3.30) imply that such a C exists for $v \geq 1$. On the other hand, an application of Cauchy's estimates in a disk of radius $|\lambda| = r < \sqrt{\pi}$ also yields

$$|r_N(\lambda)| \leq M_r \sum_{k=N}^{\infty} \left(\frac{|\lambda|}{r}\right)^k \leq M_r \left(\frac{|\lambda|}{r}\right)^N \frac{1}{1 - \frac{|\lambda|}{r}}, \qquad |\lambda| < r,$$

where $M_r = \max_{|\lambda|=r} |f(\lambda)|$. Choosing any $r \in (1, \sqrt{\pi})$ we see that such a C also exists for $0 \leq v \leq 1$. This proves (3.32) and hence also (3.30) and (3.31).

Equation (3.30) together with the identity

$$\int_0^\infty e^{-tv} v^\alpha dv = \frac{\Gamma(\alpha+1)}{t^{\alpha+1}}, \qquad \alpha > -1,$$

imply the following asymptotic expansion of $G_U^{(1)}$:

$$G_U^{(1)}(t, \sigma; t) = \left(\frac{t}{2\pi}\right)^s e^{-it} \sum_{k=0}^{N-1} \frac{c_k \Gamma(\frac{k+1}{2})}{t^{\frac{k+1}{2}}} + O\left(\frac{\Gamma(\frac{N+1}{2})}{t^{\frac{N+1}{2}-\sigma}}\right). \tag{3.33}$$

Proof of (3.20)

Equations (3.13), (3.26), and (3.33) give the asymptotics of G_L and G_U. Substitution into (3.23) yields

$$\begin{aligned}
\zeta(1-s) = & \sum_{n=1}^{[\frac{t}{2\pi}]} n^{s-1} - \frac{1}{s}\left(\frac{t}{2\pi}\right)^s \\
& + \frac{e^{\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=1}^{\infty} \sum_{j=0}^{N-1} e^{-nz + it\ln z} \left(\frac{1}{n - \frac{it}{z}} \frac{d}{dz}\right)^j \frac{z^{\sigma-1}}{n - \frac{it}{z}} \bigg|_{z=-it} \\
& + \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s} \sum_{n=2}^{\infty} \sum_{j=0}^{N-1} e^{-nz + it\ln z} \left(\frac{1}{n - \frac{it}{z}} \frac{d}{dz}\right)^j \frac{z^{\sigma-1}}{n - \frac{it}{z}} \bigg|_{z=it} \\
& + \left(\frac{t}{2\pi}\right)^s e^{-it} \sum_{k=0}^{2N} \frac{c_k(\sigma)\Gamma(\frac{k+1}{2})}{t^{\frac{k+1}{2}}} + O\left((2N+1)!!Ne^{2(N+1)}t^{\sigma-N-1}\right),
\end{aligned}$$

where we have replaced N by $2N+1$ in (3.33) and used that $\Gamma(N+1) = N! \leq (2N+1)!!$ to eliminate one of the error terms. Replacing σ by $1-\sigma$ and taking the complex conjugate of the resulting equation, we find (3.20).

Proof of (3.22)

Letting $N = 3$ in (3.20) and using the expressions in (3.21) for $\{c_k\}_0^6$, we find that the term on the rhs of (3.20) involving the c_k's yields the term involving the second curly bracket on the rhs of (3.22). On the other hand, using that

$$\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\frac{z^{-\sigma}}{n+\frac{it}{z}}=-\frac{z^{1-\sigma}(n\sigma z+i(\sigma-1)t)}{(nz+it)^3}$$

and

$$\left(\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\right)^2\frac{z^{-\sigma}}{n+\frac{it}{z}}=\frac{z^{1-\sigma}\left(n^2\sigma(\sigma+1)z^2+in\left(2\sigma^2-\sigma-2\right)tz-(\sigma-1)^2t^2\right)}{(nz+it)^5},$$

long but straightforward computations show that the two terms involving the double sums on the rhs of (3.20) yield the term involving the first curly bracket on the rhs of (3.22). This proves (3.22).

4 The Asymptotics of the Riemann Zeta Function for $0<\eta<t$

In this section, we consider the asymptotics of $\zeta(s)$ as $t\to\infty$ with $0<\eta<t$. Theorem 4.1 and its corollary treat the cases $\epsilon<\eta<\sqrt{t}$ and $2\pi\sqrt{t}<\eta<\frac{2\pi}{\epsilon}t$ under the assumption that $\mathrm{dist}(\eta,2\pi\mathbb{Z})>\epsilon$ for some $\epsilon>0$. The case when η is of the same order as $\sqrt{t}$ and the case when $\mathrm{dist}(\eta,2\pi\mathbb{Z})\to0$ are covered by theorem 4.4 and its corollary.

Theorem 4.1 (The asymptotics to all orders for the case $1<\eta<\sqrt{t}$) *For every $\epsilon>0$, there exists a constant $A>0$ such that*

$$\begin{aligned}\zeta(s)=&\sum_{n=1}^{[\frac{t}{\eta}]}\frac{1}{n^s}+\chi(s)\sum_{n=1}^{[\frac{\eta}{2\pi}]}\frac{1}{n^{1-s}} \qquad (4.1)\\ &-\frac{e^{-i\pi s}\Gamma(1-s)}{2\pi i}e^{-([\frac{t}{\eta}]+1)i\eta}e^{\frac{i\pi}{2}(s-1)}\eta^{s-1}e^{\frac{i\pi}{4}}\sum_{k=0}^{[\frac{N-1}{2}]}\frac{\varphi^{(2k)}(0)}{(2k)!}i^k\left(\frac{2\eta^2}{t}\right)^{k+\frac12}\Gamma\left(k+\frac12\right)\\ &+e^{-i\pi s}\Gamma(1-s)e^{-\frac{\pi t}{2}}\eta^{\sigma-1}\\ &\times\begin{cases}O\left(\left(\frac{3N}{t}\right)^{\frac{N}{6}}\frac{\eta}{\sqrt{t}}\right), & \epsilon<\eta<t^{\frac13}<\infty,\quad 1\le N<\frac{At}{\eta^3},\\ O\left(Ne^{-\frac{At}{\eta^2}}+\left(\frac{N\eta^2}{t}\right)^{\frac{N+1}{2}}\right), & t^{\frac13}<\eta<\sqrt{t}<\infty,\quad 1\le N<\frac{At}{\eta^2},\end{cases}\\ &\qquad\qquad \mathrm{dist}(\eta,2\pi\mathbb{Z})>\epsilon,\quad 0\le\sigma\le1,\quad t\to\infty,\end{aligned}$$

where the error terms are uniform for all η,σ,N within the above ranges, and the function $\varphi(z)$ is defined by

$$\varphi(z)=\frac{e^{(s-1)\log(1+\frac{z}{i\eta})-\frac{it}{2\eta^2}z^2-[\frac{t}{\eta}]z}}{e^z-e^{-i\eta}}.$$

For $N = 3$ equation (4.1) simplifies to

$$\begin{aligned}\zeta(s) = {} & \sum_{n=1}^{[\frac{t}{\eta}]} \frac{1}{n^s} + \chi(s) \sum_{n=1}^{[\frac{\eta}{2\pi}]} \frac{1}{n^{1-s}} \\ & - \frac{e^{-i\pi s}\Gamma(1-s)}{2\pi i} e^{-([\frac{t}{\eta}]+1)i\eta} e^{\frac{i\pi}{2}(s-1)} \eta^{s-1} e^{\frac{i\pi}{4}} \Bigg\{ \frac{\sqrt{2\pi}}{1-e^{-i\eta}} \frac{\eta}{\sqrt{t}} \\ & + \frac{1}{(1-e^{-i\eta})^3} \Bigg[\frac{(1-e^{-i\eta})^2 \big((-\eta[\frac{t}{\eta}] - i(\sigma-1) + t)^2 + \sigma - 1 \big)}{\eta^2} \\ & \qquad - \frac{2(1-e^{-i\eta})(-\eta[\frac{t}{\eta}] - i(\sigma-1) + t)}{\eta} + e^{-i\eta} + 1 \Bigg] \frac{i\sqrt{\pi}\eta^3}{\sqrt{2}t^{\frac{3}{2}}} \Bigg\} \\ & + e^{-i\pi s}\Gamma(1-s) e^{-\frac{\pi t}{2}} \eta^{\sigma-1} \times \begin{cases} O\big(\frac{\eta}{t}\big), & \epsilon < \eta < t^{\frac{1}{3}} < \infty, \quad 3\eta^3 < At, \\ O\big(e^{-\frac{At}{\eta^2}} + \frac{\eta^4}{t^2}\big), & t^{\frac{1}{3}} < \eta < \sqrt{t} < \infty, \quad 3\eta^2 < At, \end{cases} \\ & \qquad \mathrm{dist}(\eta, 2\pi\mathbb{Z}) > \epsilon, \quad 0 \le \sigma \le 1, \quad t \to \infty, \end{aligned} \tag{4.2}$$

where the error term is uniform for all η, σ in the above ranges.

Remark 4.2 It will be shown below (see equations (4.20) and (4.21)) that

$$\frac{\varphi^{(n)}(0)}{n!} = O\left(\frac{t^{\frac{n}{3}}}{\eta^n} \right) + O(1), \qquad n \ge 0, \quad \text{(not uniformly in } n).$$

Thus, the term in (4.1) which involves $\varphi^{(n)}(0)$ is of order

$$\begin{cases} O\left(e^{-i\pi s}\Gamma(1-s) e^{-\frac{\pi t}{2}} \frac{\eta^\sigma}{\sqrt{t}} \frac{1}{t^{\frac{n}{6}}} \right), & \epsilon < \eta < t^{\frac{1}{3}}, \\ O\left(e^{-i\pi s}\Gamma(1-s) e^{-\frac{\pi t}{2}} \eta^{\sigma-1} \left(\frac{2\eta^2}{t}\right)^{\frac{n+1}{2}} \right), & t^{\frac{1}{3}} < \eta < \sqrt{t}, \end{cases} \qquad \text{(not uniformly in } n).$$

It follows that, as expected, the first term left out of the sum is smaller than the error term.

Proof of theorem 4.1. We start with the expression for $\zeta(s)$ given on page 82 of Titchmarsh [9]:

$$\zeta(s) = \sum_{n=1}^{[\frac{t}{\eta}]} \frac{1}{n^s} + \chi(s) \sum_{n=1}^{[\frac{\eta}{2\pi}]} \frac{1}{n^{1-s}} + \frac{e^{-i\pi s}\Gamma(1-s)}{2\pi i} \sum_{j=1}^{4} I_j, \qquad 0 < \eta < t, \tag{4.3}$$

where

$$I_j = \int_{C_j} \frac{w^{s-1} e^{-[\frac{t}{\eta}]w}}{e^w - 1} dw,$$



Figure 5 *The critical point $i\eta$ and the integration contours $\{C_j\}_1^4$ in the complex w-plane.*

and the contours C_1, C_2, C_3, C_4 are the straight lines joining $\infty, c\eta + i\eta(1+c), -c\eta + i\eta(1-c), -c\eta - (2[\frac{\eta}{2\pi}]+1)\pi i, \infty$, where c is an absolute constant, $0 < c \le 1/2$, see figure 5. We henceforth let $c = 2^{-3/2}$ so that the total length of the contour C_2 is η. The branch cut for the logarithm now runs along the positive real axis. The contributions to $\zeta(s)$ from I_1, I_3, and I_4 are exponentially small by the arguments on pages 82-83 of [9] (in these arguments, it is assumed that $x = t/\eta \le y = \eta/2\pi$, however this assumption is not used for the derivations of these estimates and therefore we do not need to make it here.)

Let $\epsilon > 0$ be given and suppose that $2\epsilon < \eta < \sqrt{t}$, $0 \le \sigma \le 1$, $m = [t/\eta]$, $\mathrm{dist}(\eta, 2\pi\mathbb{Z}) > 2\epsilon$, and $N \ge 1$. All error terms of the form $O(\cdot)$ will be uniform with respect to η, σ, N (but not with respect to ϵ). In order to analyze the integral I_2, we write

$$I_2 = e^{-(m+1)i\eta}(i\eta)^{s-1}\int_{C_2} e^{\frac{it}{2\eta^2}(w-i\eta)^2}\varphi(w-i\eta)dw,$$

where

$$\varphi(z) = \frac{e^{(s-1)\log(1+\frac{z}{i\eta}) - \frac{t}{\eta}z - \frac{it}{2\eta^2}z^2 + (\frac{t}{\eta}-[\frac{t}{\eta}])z}}{e^z - e^{-i\eta}}.$$

Defining $\phi(z)$ by

$$\phi(z) = e^{(s-1)\log\left(1+\frac{z}{i\eta}\right) - \frac{t}{\eta}z - \frac{it}{2\eta^2}z^2}, \tag{4.4}$$

we have

$$\varphi(z) = \frac{\phi(z)e^{(\frac{t}{\eta}-[\frac{t}{\eta}])z}}{e^z - e^{-i\eta}}.$$

We write for z in a neighborhood of zero,

$$\varphi(z) = \sum_{n=0}^{\infty} b_n z^n = \sum_{n=0}^{N-1} b_n z^n + s_N(z).$$

Then

$$s_N(z) = \frac{z^N}{2\pi i}\int_\Gamma \frac{\varphi(w)dw}{w^N(w-z)}, \tag{4.5}$$

where Γ is a contour which encircles 0 and z but none of the poles of φ. We split the contour C_2 as follows:

$$C_2 = C_2^\epsilon \cup C_2^r,$$

where C_2^ϵ denotes the segment of C_2 of length 2ϵ which consists of the points within a distance ϵ from $i\eta$. We write

$$I_2 = I_2^\epsilon + I_2^r,$$

where

$$I_2^\epsilon = e^{-(m+1)i\eta}(i\eta)^{s-1}\int_{C_2^\epsilon} e^{\frac{it}{2\eta^2}(w-i\eta)^2}\varphi(w-i\eta)dw$$

and

$$I_2^r = e^{-(m+1)i\eta}(i\eta)^{s-1}\int_{C_2^r} e^{\frac{it}{2\eta^2}(w-i\eta)^2}\varphi(w-i\eta)dw.$$

We claim that

$$I_2^r = O\left(e^{-\frac{\pi t}{2}}\frac{\eta^\sigma}{\sqrt{t}}e^{-\frac{t\epsilon^2}{24\eta^2}}\right). \tag{4.6}$$

Indeed, the change of variables $w = i\eta + \lambda e^{\frac{i\pi}{4}}$ gives

$$I_2^r = O\left(e^{-\frac{\pi t}{2}}\eta^{\sigma-1}\left(\int_{-\frac{\eta}{2}}^{-\epsilon} + \int_\epsilon^{\frac{\eta}{2}}\right)e^{-\frac{t}{2\eta^2}\lambda^2}|\varphi(\lambda e^{\frac{i\pi}{4}})|d\lambda\right).$$

There exists a constant $A > 0$ such that[1]

$$\left|\frac{e^{(\frac{t}{\eta}-[\frac{t}{\eta}])z}}{e^z - e^{-i\eta}}\right| < \frac{A}{\epsilon}, \qquad z = \lambda e^{\frac{i\pi}{4}}, \quad \epsilon < |\lambda| < \frac{\eta}{2}. \tag{4.7}$$

---

[1] Here and below A denotes a strictly positive constant which may change within a computation.

Moreover, the definition of $\phi(w)$ implies

$$\begin{aligned}\log\phi(w) &= (s-1)\log\left(1+\frac{w}{i\eta}\right)-\frac{t}{\eta}w-\frac{it}{2\eta^2}w^2\\ &= (\sigma-1)\log\left(1+\frac{w}{i\eta}\right)-\frac{it}{\eta^2}w^2\sum_{k=1}^{\infty}\frac{(-1)^{k-1}}{k+2}\left(\frac{w}{i\eta}\right)^k.\end{aligned}$$

Hence, for $|w|\le\frac{3}{5}\eta$ we have

$$\operatorname{Re}\log\phi(w)\le|\sigma-1|\log\frac{8}{5}+\frac{t}{\eta^2}|w|^2\frac{1}{3}\frac{|w|}{\eta}\frac{5}{2}. \tag{4.8}$$

and so

$$|\phi(\lambda e^{\frac{i\pi}{4}})|<e^{|\sigma-1|\ln\frac{8}{5}+\frac{5t}{6\eta^3}\lambda^3},\qquad \epsilon<|\lambda|<\frac{\eta}{2}. \tag{4.9}$$

Equations (4.7) and (4.9) imply that

$$|\varphi(\lambda e^{\frac{i\pi}{4}})|=O\left(e^{\frac{5t}{6\eta^3}|\lambda|^3}\right),\qquad \epsilon<|\lambda|<\frac{\eta}{2}.$$

This yields

$$\begin{aligned}I_2^r &= O\left(e^{-\frac{\pi t}{2}}\eta^{\sigma-1}\int_{\epsilon}^{\frac{\eta}{2}}e^{-\frac{t}{2\eta^2}\lambda^2+\frac{5t}{6\eta^3}|\lambda|^3}d\lambda\right)\\ &= O\left(e^{-\frac{\pi t}{2}}\eta^{\sigma-1}\int_{\epsilon}^{\frac{\eta}{2}}e^{-\frac{t}{2\eta^2}\lambda^2+\frac{5t}{12\eta^2}\lambda^2}d\lambda\right)=O\left(e^{-\frac{\pi t}{2}}\eta^{\sigma-1}\int_{\epsilon}^{\frac{\eta}{2}}e^{-\frac{t}{12\eta^2}\lambda^2}d\lambda\right).\end{aligned}$$

Splitting the integrand as

$$e^{-\frac{t}{12\eta^2}\lambda^2}=e^{-\frac{t}{24\eta^2}\lambda^2}\times e^{-\frac{t}{24\eta^2}\lambda^2}$$

and noting that

$$\int_{\epsilon}^{\frac{\eta}{2}}e^{-\frac{t}{24\eta^2}\lambda^2}d\lambda=O\left(\frac{\eta}{\sqrt{t}}\right),$$

we find

$$I_2^r=O\left(e^{-\frac{\pi t}{2}}\eta^{\sigma-1}e^{-\frac{t}{24\eta^2}\epsilon^2}\frac{\eta}{\sqrt{t}}\right).$$

This proves (4.6).

We next claim that

$$I_2^{\epsilon}=e^{-(m+1)i\eta}(i\eta)^{s-1}\int_{C_2^{\epsilon}}e^{\frac{it}{2\eta^2}(w-i\eta)^2}\sum_{n=0}^{N-1}b_n(w-i\eta)^n dw \tag{4.10}$$

$$+\begin{cases} O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\left(\frac{3N}{t}\right)^{\frac{N}{6}}\frac{\eta}{\sqrt{t}}\right), & 1\le N<\frac{At}{\eta^3}, \\ O\left(e^{-\frac{\pi t}{2}}\eta^{\sigma-1}\left(\frac{N\eta^2}{t}\right)^{\frac{N+1}{2}}\right), & t^{\frac{1}{3}}<\eta<\sqrt{t}. \end{cases}$$

In order to establish (4.10), we need to estimate the error term

$$e^{-(m+1)i\eta}(i\eta)^{s-1}\int_{C_2^{\epsilon}} e^{\frac{it}{2\eta^2}(w-i\eta)^2}s_N(w-i\eta)dw.$$

In view of the assumption $\mathrm{dist}(\eta, 2\pi\mathbb{Z}) > 2\epsilon$, we have

$$\left|\frac{e^{(\frac{t}{\eta}-[\frac{t}{\eta}])z}}{e^z-e^{-i\eta}}\right| = O(1), \qquad |z|<\epsilon. \tag{4.11}$$

Let us first consider the case $1\le N<\frac{At}{\eta^3}$. Let $|z|<\frac{40}{21}\epsilon$ (so that $\frac{21}{20}|z|<2\epsilon$) and let Γ in (4.5) be a circle with center $w=0$ and radius ρ_N, where

$$\frac{21}{20}|z|\le\rho_N<2\epsilon.$$

Shrinking $\epsilon>0$ if necessary, we may assume that $2\epsilon<\frac{3}{5}\eta$. Then, since $\varphi(w)$ is analytic in the disk $|w|<2\epsilon$, equations (4.5), (4.8), and (4.11) yield

$$s_N(z) = O\left(\frac{2\pi\rho_N}{\rho_N-|z|}|z|^N\rho_N^{-N}e^{\frac{5t\rho_N^3}{6\eta^3}}\right) = O\left(|z|^N\rho_N^{-N}e^{\frac{5t\rho_N^3}{6\eta^3}}\right), \qquad |z|<\frac{40}{21}\epsilon.$$

The function $\rho^{-N}e^{\frac{5t\rho^3}{6\eta^3}}$ has the minimum $(\frac{5et}{2N\eta^3})^{N/3}$ for $\rho=(\frac{2N}{5t})^{1/3}\eta$; ρ_N can have this value if

$$\frac{21}{20}|z|\le\left(\frac{2N}{5t}\right)^{1/3}\eta<2\epsilon.$$

The assumption $1\le N<\frac{At}{\eta^3}$ implies that for A sufficiently small, we have $(\frac{2N}{5t})^{1/3}\eta<2\epsilon$. Hence, letting $\rho_N=(\frac{2N}{5t})^{1/3}\eta$, we find

$$s_N(z) = O\left(|z|^N\left(\frac{5et}{2N\eta^3}\right)^{\frac{N}{3}}\right), \qquad N<\frac{At}{\eta^3}, \quad |z|\le\frac{20}{21}\left(\frac{2N}{5t}\right)^{\frac{1}{3}}\eta<\frac{40}{21}\epsilon, \tag{4.12}$$

For $|z|<\frac{40}{21}\epsilon$ we can also take $\rho_N=\frac{21}{20}|z|$, which yields

$$s_N(z) = O\left(\left(\frac{20}{21}\right)^N e^{\frac{5}{6}\frac{t}{\eta^3}(\frac{21}{20}|z|)^3}\right) = O\left(e^{\frac{14}{29}\frac{t}{\eta^2}|z|^2}\right), \qquad |z|\le\epsilon. \tag{4.13}$$

Using (4.12) and (4.13), we estimate

$$e^{-(m+1)i\eta}(i\eta)^{s-1}\int_{C_2^{\epsilon}} e^{\frac{it}{2\eta^2}(w-i\eta)^2}s_N(w-i\eta)dw \tag{4.14}$$

$$
\begin{aligned}
&= O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\biggl\{\int_0^{A(\frac{N}{t})^{\frac{1}{3}}\eta} e^{-\frac{t}{2\eta^2}\lambda^2}\lambda^N\left(\frac{5et}{2N\eta^3}\right)^{\frac{N}{3}}d\lambda + \int_{A(\frac{N}{t})^{\frac{1}{3}}\eta}^{\epsilon} e^{-\frac{t}{2\eta^2}\lambda^2+\frac{14}{29}\frac{t}{\eta^2}\lambda^2}d\lambda\biggr\}\biggr)\\
&= O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\biggl\{\left(\frac{5et}{2N\eta^3}\right)^{\frac{N}{3}}2^{\frac{N-1}{2}}\left(\frac{\eta^2}{t}\right)^{\frac{N+1}{2}}\Gamma\left(\frac{N+1}{2}\right) + \int_{A(\frac{N}{t})^{\frac{1}{3}}\eta}^{\epsilon} e^{-\frac{t}{58\eta^2}\lambda^2}d\lambda\biggr\}\biggr).\\
&= O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\left(\frac{5et}{2N\eta^3}\right)^{\frac{N}{3}}2^{\frac{N-1}{2}}\left(\frac{\eta^2}{t}\right)^{\frac{N+1}{2}}\Gamma\left(\frac{N+1}{2}\right)\biggr), \qquad 1 \le N < \frac{At}{\eta^3}.
\end{aligned}
$$

where we have used the following estimate to find the last equality:

$$
\begin{aligned}
\int_{A(\frac{N}{t})^{\frac{1}{3}}\eta}^{\epsilon} e^{-\frac{t}{58\eta^2}\lambda^2}d\lambda &= O\biggl(e^{-\frac{t}{116\eta^2}(A(\frac{N}{t})^{\frac{1}{3}}\eta)^2}\int_{A(\frac{N}{t})^{\frac{1}{3}}\eta}^{\infty} e^{-\frac{t}{116\eta^2}\lambda^2}d\lambda\biggr)\\
&= O\biggl(e^{-\frac{A^2t^{\frac{1}{3}}}{116}N^{\frac{2}{3}}}\frac{\eta}{\sqrt{t}}\biggr) = O(e^{-\frac{A^2}{116}t^{\frac{1}{3}}}).
\end{aligned}
$$

Using the expansion

$$
\Gamma(x) = e^{x(\log(x)-1)}\biggl(\sqrt{\frac{2\pi}{x}} + O\biggl(\frac{1}{x^{3/2}}\biggr)\biggr), \qquad x \to \infty,
$$

and the inequality $\frac{25}{4e} < 3$, it follows that the right-hand side of (4.14) is

$$
O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\left(\frac{3N}{t}\right)^{\frac{N}{6}}\frac{\eta}{\sqrt{t}}\biggr). \tag{4.15}
$$

This proves (4.10) in the case when $1 \le N < \frac{At}{\eta^3}$.

We now consider the case when $t^{\frac{1}{3}} < \eta < \sqrt{t}$. Using (4.11) and (4.8) in the representation (4.5) with Γ a circle with center $w = 0$ and radius $2^{1/3}\epsilon$, we find that

$$
s_N(z) = O\biggl(\frac{|z|^N}{\epsilon^{N+1}}e^{\frac{5t\epsilon^3}{3\eta^3}}\biggr), \qquad |z| < \epsilon. \tag{4.16}
$$

Now $t^{\frac{1}{3}} < \eta$ implies that $e^{\frac{5t\epsilon^3}{3\eta^3}} = O(1)$, so we can estimate

$$
\begin{aligned}
&e^{-(m+1)i\eta}(i\eta)^{s-1}\int_{C_2^\epsilon} e^{\frac{it}{2\eta^2}(w-i\eta)^2}s_N(w-i\eta)dw\\
&\qquad = O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\int_0^{\epsilon} e^{-\frac{t}{2\eta^2}\lambda^2}\lambda^N d\lambda\biggr)\\
&\qquad = O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}2^{\frac{N-1}{2}}\left(\frac{\eta^2}{t}\right)^{\frac{N+1}{2}}\Gamma\left(\frac{N+1}{2}\right)\biggr)\\
&\qquad = O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\left(\frac{\eta^2}{t}\right)^{\frac{N+1}{2}}N^{\frac{N}{2}}\biggr), \qquad t^{\frac{1}{3}} < \eta < \sqrt{t}.
\end{aligned}
$$

This completes the proof of (4.10).

We next claim that, up to a small error term, the contour C_2^ϵ in the integral in (4.10) can be replaced by the infinite line C_2', where C_2' denotes the infinite straight line of which C_2 is a part. More precisely, we claim that there exists an $A > 0$ such that

$$e^{-(m+1)i\eta}(i\eta)^{s-1}\int_{C_2'\setminus C_2^\epsilon} e^{\frac{it}{2\eta^2}(w-i\eta)^2}\sum_{n=0}^{N-1} b_n(w-i\eta)^n dw \tag{4.17}$$
$$= \begin{cases} O\Big(e^{-\frac{\pi t}{2}-\frac{At}{\eta^2}}\Big), & 2\epsilon < \eta < t^{\frac{1}{3}}, \quad 1 \le N < \frac{At}{\eta^3}, \\ O\Big(\eta^{\sigma-1}e^{-\frac{\pi t}{2}-\frac{At}{\eta^2}}N\Big), & t^{\frac{1}{3}} < \eta < \sqrt{t}, \quad 1 \le N < \frac{At}{\eta^2}. \end{cases}$$

In order to prove (4.17), we note that the coefficient of b_n on the lhs of (4.17) is

$$O\Big(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\int_\epsilon^\infty e^{-\frac{t}{2\eta^2}\lambda^2}\lambda^n d\lambda\Big). \tag{4.18}$$

We write the integrand as

$$e^{-\frac{t}{4\eta^2}\lambda^2} \times e^{-\frac{t}{4\eta^2}\lambda^2}\lambda^n,$$

and the second factor is steadily decreasing for $\lambda > \sqrt{\frac{2n}{t}}\eta$, and so it decreases throughout the interval of integration provided that $n < N < \frac{t\epsilon^2}{2\eta^2}$. The term in (4.18) is then

$$O\Big(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}e^{-\frac{t}{4\eta^2}\epsilon^2}\epsilon^n\int_\epsilon^\infty e^{-\frac{t}{4\eta^2}\lambda^2}d\lambda\Big) = O\Big(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}e^{-\frac{t}{4\eta^2}\epsilon^2}\epsilon^n\frac{\eta}{\sqrt{t}}\Big). \tag{4.19}$$

We now assume that $N < At/\eta^3$. Choosing $|z| < \frac{20}{21}\big(\frac{2}{5t}\big)^{\frac{1}{3}}\eta$, equation (4.12) yields

$$b_n = (s_n(z) - s_{n+1}(z))z^{-n} = O\left(\left(\frac{5et}{2n\eta^3}\right)^{\frac{n}{3}}\right), \qquad N < \frac{At}{\eta^3}, \quad n \ge 1. \tag{4.20}$$

Multiplying the rhs of (4.19) by b_n and summing from 0 to $N-1$, we find that the total error is

$$O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}e^{-\frac{t}{4\eta^2}\epsilon^2}\frac{\eta}{\sqrt{t}}\Big(b_0 + \sum_{n=1}^{N-1}\epsilon^n\left(\frac{5et}{2n\eta^3}\right)^{\frac{n}{3}}\Big)\right).$$

Now the function $(\frac{t}{n\eta^3})^{\frac{n}{3}}$ increases steadily up to $n = \frac{t}{\eta^3 e}$, so that if $n < A\frac{t}{\eta^3}$, where $A < 1/e$, it is of order

$$O\Big(e^{\frac{1}{3}\frac{At}{\eta^3}\ln\frac{1}{A}}\Big).$$

Hence, if $\epsilon^3 < \frac{2}{5e}$ and $N < A\frac{t}{\eta^3}$, with A small enough that $\frac{1}{3\epsilon}A\ln\frac{1}{A} < \frac{\epsilon^2}{4}$, the total error is

$$O\Big(e^{-\frac{\pi t}{2}-\frac{At}{\eta^2}}\Big).$$

This proves (4.17) when $N < A\frac{t}{\eta^3}$. In the case when $t^{\frac{1}{3}} < \eta < \sqrt{t}$, we instead use (4.16) to find

$$b_n = (s_n(z) - s_{n+1}(z))z^{-n} = O\Big(e^{\frac{5t\epsilon^3}{3\eta^3}}\Big), \qquad n \geq 0. \tag{4.21}$$

Then, since $t^{\frac{1}{3}} < \eta$, the total error is

$$O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}-\frac{t}{4\eta^2}\epsilon^2}e^{\frac{5t\epsilon^3}{3\eta^3}}N\right) = O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}-\frac{At}{\eta^2}}N\right), \qquad t^{\frac{1}{3}} < \eta < \sqrt{t}.$$

This completes the proof of (4.17).

We finally analyze the sum

$$\begin{aligned}
&e^{-(m+1)i\eta}(i\eta)^{s-1}\int_{C_2'} e^{\frac{it}{2\eta^2}(w-i\eta)^2}\sum_{n=0}^{N-1} b_n(w-i\eta)^n dw \\
&= -e^{-(m+1)i\eta}(i\eta)^{s-1}\sum_{n=0}^{N-1} b_n e^{\frac{\pi i}{4}(n+1)}\int_{-\infty}^{\infty} e^{-\frac{t}{2\eta^2}\lambda^2}\lambda^n d\lambda \\
&= -e^{-(m+1)i\eta}(i\eta)^{s-1}\sum_{n=0}^{N-1} b_n e^{\frac{\pi i}{4}(n+1)}2^{\frac{n-1}{2}}(1+(-1)^n)\Big(\frac{\eta^2}{t}\Big)^{\frac{n+1}{2}}\Gamma\Big(\frac{n+1}{2}\Big) \\
&= -e^{-(m+1)i\eta}(i\eta)^{s-1}e^{\frac{i\pi}{4}}\sum_{k=0}^{[\frac{N-1}{2}]} b_{2k}i^k\Big(\frac{2\eta^2}{t}\Big)^{k+\frac{1}{2}}\Gamma\Big(k+\frac{1}{2}\Big).
\end{aligned}$$

Together with equations (4.3), (4.6), (4.10), and (4.17), this yields (4.1). **QED**

Corollary 4.3 (The asymptotics to all orders for the case $2\pi\sqrt{t} < \eta < \frac{2\pi}{\epsilon}t$) *For every $\epsilon > 0$, there exists a constant $A > 0$ such that*

$$\begin{aligned}
\zeta(s) = &\sum_{n=1}^{[\frac{t}{\eta}]}\frac{1}{n^s} + \chi(s)\sum_{n=1}^{[\frac{\eta}{2\pi}]}\frac{1}{n^{1-s}} \\
&+ \chi(s)\frac{e^{i\pi(1-s)}\Gamma(s)}{2\pi i}e^{([\frac{\eta}{2\pi}]+1)i\frac{2\pi t}{\eta}}e^{\frac{i\pi s}{2}}\Big(\frac{2\pi t}{\eta}\Big)^{-s}e^{-\frac{i\pi}{4}}\sum_{k=0}^{[\frac{N-1}{2}]}\frac{\psi^{(2k)}(0)}{(2k)!}(-i)^k\Big(\frac{8\pi^2 t}{\eta^2}\Big)^{k+\frac{1}{2}}\Gamma\Big(k+\frac{1}{2}\Big) \\
&+ \chi(s)e^{i\pi(1-s)}\Gamma(s)e^{-\frac{\pi t}{2}}\Big(\frac{\eta}{2\pi t}\Big)^{\sigma} \\
&\times\begin{cases} O\Big(\big(\frac{3N}{t}\big)^{\frac{N}{6}}\frac{\frac{2\pi t}{\eta}}{\sqrt{t}}\Big), & 2\pi t^{\frac{2}{3}} < \eta < \frac{2\pi}{\epsilon}t < \infty, \quad 1 \leq N < \frac{A\eta^3}{t^2}, \\ O\Big(Ne^{-\frac{A\eta^2}{4\pi^2 t}} + \big(\frac{N4\pi^2 t}{\eta^2}\big)^{\frac{N+1}{2}}\Big), & 2\pi\sqrt{t} < \eta < 2\pi t^{\frac{2}{3}} < \infty, \quad 1 \leq N < \frac{A\eta^2}{t}, \end{cases}
\end{aligned} \tag{4.22}$$

$$\mathrm{dist}\left(\frac{2\pi t}{\eta}, 2\pi\mathbb{Z}\right) > \epsilon, \quad 0 \le \sigma \le 1, \quad t \to \infty,$$

where the error terms are uniform for all η, σ, N within the above ranges, and the function $\psi(z)$ is defined by

$$\psi(z) = \frac{e^{-s\log(1+\frac{i\eta z}{2\pi t})+\frac{i\eta^2}{8\pi^2 t}z^2-[\frac{\eta}{2\pi}]z}}{e^z - e^{\frac{2\pi i t}{\eta}}}.$$

Proof. We replace σ by $1-\sigma$ in (4.1) and take the complex conjugate of both sides. We then multiply the resulting equation by $\chi(s)$ and use the identities

$$\begin{aligned} &\overline{\zeta(\bar{s})} = \zeta(s), \qquad \overline{\Gamma(\bar{s})} = \Gamma(s), \qquad \overline{\chi(\bar{s})} = \chi(s), \\ &\chi(s)\chi(1-s) = 1, \qquad \chi(s)\zeta(1-s) = \zeta(s). \end{aligned} \tag{4.23}$$

This yields the following equation

$$\begin{aligned} \zeta(s) = {}& \chi(s)\sum_{n=1}^{[\frac{t}{\eta}]}\frac{1}{n^{1-s}} + \sum_{n=1}^{[\frac{\eta}{2\pi}]}\frac{1}{n^s} \qquad\qquad (4.24)\\ &+ \chi(s)\frac{e^{i\pi(1-s)}\Gamma(s)}{2\pi i}e^{([\frac{t}{\eta}]+1)i\eta}e^{\frac{i\pi s}{2}}\eta^{-s}e^{-\frac{i\pi}{4}} \\ &\times \sum_{k=0}^{[\frac{N-1}{2}]}\left(\frac{d^{2k}}{dz^{2k}}\bigg|_{z=0}\frac{e^{-s\log(1-\frac{z}{i\eta})+\frac{it}{2\eta^2}z^2-[\frac{t}{\eta}]z}}{e^z - e^{i\eta}}\right)\frac{1}{(2k)!}(-i)^k\left(\frac{2\eta^2}{t}\right)^{k+\frac{1}{2}}\Gamma\left(k+\frac{1}{2}\right) \\ &+ \chi(s)e^{i\pi(1-s)}\Gamma(s)e^{-\frac{\pi t}{2}}\eta^{-\sigma} \\ &\times \begin{cases} O\left(\left(\frac{3N}{t}\right)^{\frac{N}{6}}\frac{\eta}{\sqrt{t}}\right), & \epsilon < \eta < t^{\frac{1}{3}} < \infty, \quad 1 \le N < \frac{At}{\eta^3}, \\ O\left(Ne^{-\frac{At}{\eta^2}} + \left(\frac{N\eta^2}{t}\right)^{\frac{N+1}{2}}\right), & t^{\frac{1}{3}} < \eta < \sqrt{t} < \infty, \quad 1 \le N < \frac{At}{\eta^2}, \end{cases} \\ &\qquad\qquad \mathrm{dist}(\eta, 2\pi\mathbb{Z}) > \epsilon, \quad 0 \le \sigma \le 1, \quad t \to \infty, \end{aligned}$$

Replacing η by $\frac{2\pi t}{\eta}$, we find (4.22). **QED**

Theorem 4.4 (The asymptotics to all orders for the case $\epsilon\sqrt{t} < \eta < t$) *For every $\epsilon > 0$, there exists a constant $A > 0$ such that*

$$\begin{aligned} \zeta(s) = {}& \sum_{n=1}^{[\frac{t}{\eta}]}\frac{1}{n^s} + \chi(s)\sum_{n=1}^{[\frac{\eta}{2\pi}]}\frac{1}{n^{1-s}} \qquad\qquad (4.25)\\ &+ e^{-i\pi s}\Gamma(1-s)\Bigg\{e^{\frac{i\pi(s-1)}{2}}\eta^{s-1}e^{\frac{2t}{\eta}[\frac{\eta}{2\pi}]\pi i - it - \frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi-\eta)^2}S_N(s,\eta) \end{aligned}$$

$$+\,O\left(e^{-\frac{\pi t}{2}}\left(\frac{3N}{t}\right)^{\frac{N}{6}}\frac{\eta^{\sigma}}{\sqrt{t}}\right)\Bigg\},$$

$$\epsilon\sqrt{t} < \eta < t, \quad 0 \le \sigma \le 1, \quad 1 \le N < At, \quad t \to \infty,$$

where the error term is uniform for all η, σ, N *in the above ranges and the function* $S_N(s,\eta)$ *is defined by*

$$S_N(s,\eta) = \sum_{n=0}^{N-1} a_n \sum_{k=0}^{n} \binom{n}{k} \left(2\left[\frac{\eta}{2\pi}\right]\pi i - i\eta\right)^{k} \times \partial_2^{n-k}\Phi\left(-\frac{2\pi t}{\eta^2}, \frac{2t}{\eta} - \frac{2\pi t}{\eta^2}\left[\frac{\eta}{2\pi}\right] - \left[\frac{t}{\eta}\right] - \frac{1}{2}\right). \tag{4.26}$$

The coefficients a_n *are determined by the recurrence formula*

$$i\eta(n+1)a_{n+1} = (\sigma - n - 1)a_n - \frac{it}{\eta^2}a_{n-2}, \qquad n = 0, 1, 2, \ldots, \tag{4.27}$$

together with the initial conditions $a_{-2} = a_{-1} = 0$ *and* $a_0 = 1$, *and the function* $\Phi(\tau, u)$ *is defined by*

$$\Phi(\tau, u) = \int_{0\nwarrow 1} \frac{e^{\pi i \tau x^2 + 2\pi i u x}}{e^{\pi i x} - e^{-\pi i x}}dx, \qquad \tau < 0, \quad u \in \mathbb{C}, \tag{4.28}$$

with the contour $0 \nwarrow 1$ *denoting a straight line parallel to* $e^{3\pi i/4}$ *which crosses the real axis between* 0 *and* 1.

In the particular case when $p, q > 0$ *are integers, we have*

$$\Phi\left(-\frac{p}{q}, u\right) = \frac{1}{1 - (-1)^q e^{-\pi i q p - 2\pi i q u}} \times \left\{\sum_{n=0}^{q-1}(-1)^n e^{-\pi i n^2 \frac{p}{q} - 2\pi i n u} + \frac{(-1)^q e^{-\pi i q p - 2\pi i q u}}{\sqrt{p/q}} e^{\frac{3\pi i}{4}} \sum_{n=0}^{p-1} e^{\frac{\pi i q}{p}(u + n + \frac{1}{2})^2}\right\}. \tag{4.29}$$

For $N = 3$ *equation (4.25) simplifies to*

$$\zeta(s) = \sum_{n=1}^{[\frac{t}{\eta}]} \frac{1}{n^s} + \chi(s) \sum_{n=1}^{[\frac{\eta}{2\pi}]} \frac{1}{n^{1-s}} \tag{4.30}$$

$$+\, e^{-i\pi s}\Gamma(1-s)\Bigg\{ e^{\frac{i\pi(s-1)}{2}} \eta^{s-1} e^{\frac{2t}{\eta}[\frac{\eta}{2\pi}]\pi i - it - \frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi - \eta)^2}$$

$$\times \Bigg[\Phi + \frac{\sigma - 1}{i\eta}\left(\partial_2\Phi + \left(2\left[\frac{\eta}{2\pi}\right]\pi i - i\eta\right)\Phi\right)$$

$$- \frac{(\sigma-2)(\sigma-1)}{2\eta^2}\left(\partial_2^2\Phi + 2\left(2\left[\frac{\eta}{2\pi}\right]\pi i - i\eta\right)\partial_2\Phi + \left(2\left[\frac{\eta}{2\pi}\right]\pi i - i\eta\right)^2\Phi\right)\Bigg]$$

$$+O\left(e^{-\frac{\pi t}{2}}\frac{\eta^{\sigma}}{t}\right)\Bigg\}, \qquad \epsilon\sqrt{t} < \eta < t, \quad 0 \le \sigma \le 1, \quad t \to \infty,$$

where the error term is uniform for all η, σin the above ranges and Φ and its partial derivatives are evaluated at the point (1.8).

Remark 4.5 1. If $\eta = \sqrt{\frac{2\pi t}{b}}$ where $b = p/q > 0$ is a rational number, then equations (4.25)-(4.29) provide an explicit asymptotic expansion of $\zeta(s)$ to all orders. The particular case $b = 1$ is the famous case analyzed by Siegel in [8] using ideas from Riemann's unpublished notes. We note that, building on the work of [5], an elementary (but formal) derivation of the Riemann-Siegel formula is presented in [2].

2. The results of theorem 4.4 are convenient for studying the higher-order asymptotics of $\zeta(s)$ in the case when $\eta = \text{constant} \times \sqrt{t}$. However, if the order of η is strictly smaller or larger than the order of $\sqrt{t}$, the results of theorem 4.4 are less convenient, because in this case the asymptotics of $\zeta(s)$ involves the asymptotics of the sums in (4.29) as p and/or q tend to infinity. In this case, the alternative representation of the asymptotics of theorem 4.1, which avoids the appearance of the above sums, is usually more convenient.

3. The error term in (4.25) is uniform with respect to η, σ, N. More precisely, this means that equation (4.25) is equivalent to the following statement: For every $\epsilon > 0$, there exist constants $A > 0$, $C > 0$, $K > 0$ such that the inequality

$$\Bigg|\zeta(s) - \sum_{n=1}^{[\frac{t}{\eta}]}\frac{1}{n^s} - \chi(s)\sum_{n=1}^{[\frac{\eta}{2\pi}]}\frac{1}{n^{1-s}}$$
$$- e^{-i\pi s}\Gamma(1-s)e^{\frac{i\pi(s-1)}{2}}\eta^{s-1}e^{\frac{2t}{\eta}[\frac{\eta}{2\pi}]\pi i - it - \frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi-\eta)^2}S_N(s,\eta)\Bigg|$$
$$\le Ce^{-i\pi s}\Gamma(1-s)e^{-\frac{\pi t}{2}}\left(\frac{3N}{t}\right)^{\frac{N}{6}}\frac{\eta^{\sigma}}{\sqrt{t}},$$

holds for all $t > K$ and all σ, N, η such that $0 \le \sigma \le 1$, $1 \le N < At$, $\epsilon\sqrt{t} < \eta < t$.

4. The main difference between the proofs of theorems 4.1 and 4.4 can be explained as follows. Consider the representation (4.3) of $\zeta(s)$. If η is of strictly smaller order than $t^{1/2}$, the main contribution to the asymptotics of $\zeta(s)$ comes from the part of the contour C_2 that lies within a distance ϵ of the critical point $i\eta$; the remaining part of C_2 gives a contribution which is suppressed by a factor of the form $e^{-\text{const}\times\frac{t}{\eta^2}}$ cf. Eq. (4.6). Thus, the proof of theorem 4.1 relies on a direct study of the integral near the critical point $i\eta$ using the method of steepest descent. On the other hand, in the case when $\eta \sim \sqrt{t}$, the asymptotic expansion of $\zeta(s)$ depends on the integral along all of the contour C_2. The contour C_2 is a line segment of total length η centered on the critical point $i\eta$. In the proof of theorem 4.4, we handle this integral by relating it to the function $\Phi(\tau, u)$ defined in (4.28).

5. The coefficients a_n satisfy the estimate

$$a_n = O\left(\frac{t^{[\frac{n}{3}]}}{\eta^n}\right), \qquad \text{(not uniformly in } n\text{)};$$

indeed, assuming this estimate up to n, we find

$$a_{n+1} = O\left(\frac{t^{[\frac{n}{3}]}}{\eta^{n+1}}\right) + O\left(\frac{t}{\eta^3}\frac{t^{[\frac{n-2}{3}]}}{\eta^{n-2}}\right) = O\left(\frac{t^{[\frac{n+1}{3}]}}{\eta^{n+1}}\right), \qquad \text{(not uniformly in } n\text{)}.$$

Proof of theorem 4.4. Let $t > 0$, $0 \leq \sigma \leq 1$, and $m = [t/\eta]$. As in the proof of theorem 4.1, $\zeta(s)$ is given by (4.3) where the contributions from I_1, I_3, and I_4 are exponentially small, and the error terms of the form $O(\cdot)$ are uniform with respect to the variables η, σ, N as $t \to \infty$. It remains to analyze the integral I_2. In the neighborhood of $w = i\eta$ we have

$$\begin{aligned}(s-1)\log\frac{w}{i\eta} &= (s-1)\left(\frac{w-i\eta}{i\eta} - \frac{1}{2}\left(\frac{w-i\eta}{i\eta}\right)^2 + \cdots\right)\\ &= \frac{t}{\eta}(w-i\eta) + \frac{it}{2\eta^2}(w-i\eta)^2 + \cdots.\end{aligned}$$

Hence we write

$$e^{(s-1)\log\frac{w}{i\eta}} = e^{\frac{t}{\eta}(w-i\eta)+\frac{it}{2\eta^2}(w-i\eta)^2}\phi(w-i\eta),$$

where $\phi(z)$ is defined by (4.4). Define $\{a_n\}_0^\infty$ by

$$\phi(z) = \sum_{n=0}^{\infty} a_n z^n.$$

The identity

$$\frac{d\phi}{dz} = \left(\frac{s-1}{i\eta+z} - \frac{t}{\eta} - \frac{it}{\eta^2}z\right)\phi(z)$$

implies

$$(i\eta+z)\sum_{n=1}^{\infty} n a_n z^{n-1} = \left[s-1-(i\eta+z)\left(\frac{t}{\eta}+\frac{itz}{\eta^2}\right)\right]\sum_{n=0}^{\infty} a_n z^n.$$

Hence the coefficients a_n are determined in succession by the recurrence formula (4.27) supplemented with the conditions $a_{-2} = a_{-1} = 0$ and $a_0 = 1$. The first few coefficients are given by

$$\begin{aligned}&a_0 = 1, \qquad a_1 = \frac{\sigma-1}{i\eta}, \qquad a_2 = -\frac{(\sigma-2)(\sigma-1)}{2\eta^2},\\ &a_3 = \frac{-2t + i(\sigma-3)(\sigma-2)(\sigma-1)}{6\eta^3},\end{aligned}$$

$$a_4 = \frac{(\sigma-4)(\sigma-3)(\sigma-2)(\sigma-1) + 2i(4\sigma-7)t}{24\eta^4}.$$

Representing $\phi(z)$ in the form

$$\phi(z) = \sum_{n=0}^{N-1} a_n z^n + r_N(z),$$

we find

$$r_N(z) = \frac{1}{2\pi i}\int_\Gamma \frac{\phi(w) z^N}{w^N(w-z)}dw, \qquad |z| < \eta,$$

where Γ is a contour contained in the disk of radius η centered at the origin which encircles the points 0 and z once.

Let $|z| < \frac{4}{7}\eta$ (so that $\frac{21}{20}|z| < \frac{3}{5}\eta$) and let Γ be a circle with center $w = 0$ and radius ρ_N, where

$$\frac{21}{20}|z| \le \rho_N \le \frac{3}{5}\eta.$$

Then, by (4.8),

$$r_N(z) = O\left(\frac{2\pi\rho_N}{\rho_N - |z|}|z|^N \rho_N^{-N} e^{\frac{5t\rho_N^3}{6\eta^3}}\right) = O\left(|z|^N \rho_N^{-N} e^{\frac{5t\rho_N^3}{6\eta^3}}\right), \qquad |z| < \frac{4}{7}\eta.$$

The function $\rho^{-N} e^{\frac{5t\rho^3}{6\eta^3}}$ has the minimum $(\frac{5et}{2N\eta^3})^{N/3}$ for $\rho = (\frac{2N}{5t})^{1/3}\eta$; ρ_N can have this value if

$$\frac{21}{20}|z| \le \left(\frac{2N}{5t}\right)^{\frac{1}{3}}\eta \le \frac{3}{5}\eta.$$

Hence,

$$r_N(z) = O\left(|z|^N \left(\frac{5et}{2N\eta^3}\right)^{\frac{N}{3}}\right), \qquad N \le \frac{27}{50}t, \quad |z| \le \frac{20}{21}\left(\frac{2N}{5t}\right)^{\frac{1}{3}}\eta. \tag{4.31}$$

For $|z| \le \frac{4}{7}\eta$ we can also take $\rho_N = \frac{21}{20}|z|$, which yields

$$r_N(z) = O\left(\left(\frac{20}{21}\right)^N e^{\frac{5}{6}\frac{t}{\eta^3}(\frac{21}{20}|z|)^3}\right) = O\left(e^{\frac{14}{29}\frac{t}{\eta^2}|z|^2}\right), \qquad |z| \le \frac{\eta}{2}. \tag{4.32}$$

We write I_2 in the form

$$I_2 = I_2^S + I_2^R, \tag{4.33}$$

where I_2^S and I_2^R denote the integrals

$$I_2^S = \int_{C_2} (i\eta)^{s-1} \frac{e^{\frac{t}{\eta}(w-i\eta) + \frac{it}{2\eta^2}(w-i\eta)^2 - mw}}{e^w - 1} \sum_{n=0}^{N-1} a_n (w-i\eta)^n dw \tag{4.34}$$

and

$$I_2^R = \int_{C_2} (i\eta)^{s-1} \frac{e^{\frac{t}{\eta}(w-i\eta)+\frac{it}{2\eta^2}(w-i\eta)^2-mw}}{e^w-1} r_N(w-i\eta)dw. \tag{4.35}$$

We claim that

$$I_2^R = O\biggl(e^{-\frac{\pi t}{2}} \left(\frac{3N}{t}\right)^{\frac{N}{6}} \frac{\eta^\sigma}{\sqrt{t}}\biggr). \tag{4.36}$$

Indeed, suppose first that $|e^w - 1| > A$ on C_2. Then we have for $w \in C_2$, $|w - i\eta| < \epsilon$, the estimate

$$\left|\frac{e^{(\frac{t}{\eta}-m)w}}{e^w-1}\right| < \frac{e^\epsilon}{A},$$

whereas for $w \in C_2$, $|w - i\eta| > \epsilon$, we have the estimates

$$\frac{e^{(\frac{t}{\eta}-m)w}}{e^w-1} = \begin{cases} O\biggl(\frac{e^{(\frac{t}{\eta}-m-1)\operatorname{Re} w}}{1-e^{-\operatorname{Re} w}}\biggr), & \operatorname{Re} w \geq 0, \\ O\biggl(\frac{e^{(\frac{t}{\eta}-m)\operatorname{Re} w}}{e^{\operatorname{Re} w}-1}\biggr), & \operatorname{Re} w < 0. \end{cases} \tag{4.37}$$

It follows from these estimates that the factor $\frac{e^{(\frac{t}{\eta}-m)w}}{e^w-1}$ in the integrand of I_2^R is of $O(1)$ on C_2. We now make the change of variables $w = i\eta + \lambda e^{\frac{i\pi}{4}}$ and split the integral into two integrals; in the first integral $|\lambda| \leq A(\frac{N}{t})^{\frac{1}{3}}\eta$, while in the second integral $A(\frac{N}{t})^{\frac{1}{3}}\eta \leq |\lambda| \leq \eta/2$. In the first integral, we use the estimate (4.31) of r_N, whereas in the second integral, we use the estimate (4.32). This yields

$$I_2^R = O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\biggl\{\int_0^{A(\frac{N}{t})^{\frac{1}{3}}\eta} e^{-\frac{t}{2\eta^2}\lambda^2}\lambda^N\left(\frac{5et}{2N\eta^3}\right)^{\frac{N}{3}}d\lambda + \int_{A(\frac{N}{t})^{\frac{1}{3}}\eta}^{\frac{\eta}{2}} e^{-\frac{t}{2\eta^2}\lambda^2+\frac{14}{29}\frac{t}{\eta^2}\lambda^2}d\lambda\biggr\}\biggr).$$

Steps almost identical to those leading from (4.14) to (4.15) now show that

$$I_2^R = O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\left(\frac{3N}{t}\right)^{\frac{N}{6}}\frac{\eta}{\sqrt{t}}\biggr).$$

The case where the contour goes near a pole gives a similar result. Indeed, if the contour passes near the pole at $w = 2w\pi i$, say within a distance 1/2 of it, we take it around an arc of the circle $|w - i\eta| = 1$. Letting $w = i\eta + e^{i\theta}$ and using (4.31), the total contribution to I_2^R from this arc is given by

$$O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\int_0^{2\pi}\left|e^{(\frac{t}{\eta}-[\frac{t}{\eta}])e^{i\theta}+\frac{it}{2\eta^2}e^{2i\theta}}r_N(e^{i\theta})\right|d\theta\biggr)$$
$$= O\biggl(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\left(\frac{5et}{2N\eta^3}\right)^{\frac{N}{3}}\int_0^{2\pi}e^{-\frac{t}{2\eta^2}\sin(2\theta)}d\theta\biggr)$$

$$= O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\left(\frac{5et}{2N\eta^3}\right)^{\frac{N}{3}}\right) = O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\left(\frac{3N}{t}\right)^{\frac{N}{6}}\frac{\eta}{\sqrt{t}}\right),$$

where we used the assumption that $\eta > \epsilon\sqrt{t}$. This proves (4.36).

We next consider I_2^S. We claim that there exists a constant $A > 0$ such that

$$I_2^S = \int_{C_2'} (i\eta)^{s-1}\frac{e^{\frac{t}{\eta}(w-i\eta)+\frac{i}{2}\frac{t}{\eta^2}(w-i\eta)^2-mw}}{e^w-1}\sum_{n=0}^{N-1} a_n(w-i\eta)^n dw + O(e^{-\frac{\pi t}{2}-At}), \qquad (4.38)$$

where C_2' denotes the infinite straight line of which C_2 is a part. Indeed, in view of (4.37), if we replace C_2 by C_2', the integral multiplying a_n in the expression for I_2^S changes by

$$O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}\int_{\frac{\eta}{2}}^{\infty} e^{-\frac{t}{2\eta^2}\lambda^2}\lambda^n d\lambda\right). \qquad (4.39)$$

We can write the integrand as

$$e^{-\frac{t}{4\eta^2}\lambda^2} \times e^{-\frac{t}{4\eta^2}\lambda^2}\lambda^n,$$

and the second factor is steadily decreasing for $\lambda > \sqrt{\frac{2n}{t}}\eta$, and so it decreases throughout the interval of integration provided that $n < N < At$, with A sufficiently small. The term in (4.39) is then

$$O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}e^{-\frac{t}{4\eta^2}\frac{\eta^2}{4}}\left(\frac{\eta}{2}\right)^n\int_{\frac{\eta}{2}}^{\infty} e^{-\frac{t}{4\eta^2}\lambda^2}d\lambda\right) = O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}}e^{-\frac{t}{4\eta^2}\frac{\eta^2}{4}}\left(\frac{\eta}{2}\right)^n\frac{\eta}{\sqrt{t}}\right). \qquad (4.40)$$

Also, by (4.31), choosing z such that $|z| \le \frac{20}{21}(\frac{2}{5t})^{\frac{1}{3}}\eta$, we find

$$\begin{aligned} a_n &= (r_n(z) - r_{n+1}(z))z^{-n} \\ &= O\left(\left(|z|^n\left(\frac{5et}{2n\eta^3}\right)^{\frac{n}{3}} - |z|^{n+1}\left(\frac{5et}{2(n+1)\eta^3}\right)^{\frac{n+1}{3}}\right)|z|^{-n}\right) \\ &= O\left(\left(\frac{5et}{2n\eta^3}\right)^{\frac{n}{3}}\right), \qquad n \ge 1. \end{aligned} \qquad (4.41)$$

Multiplying (4.40) by a_n and summing from 0 to $N-1$, we find that the total error is

$$\begin{aligned} &O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}-\frac{t}{16}}\frac{\eta}{\sqrt{t}}\left(a_0 + \sum_{n=1}^{N-1}\left(\frac{\eta}{2}\right)^n\left(\frac{5et}{2n\eta^3}\right)^{\frac{n}{3}}\right)\right) \\ &= O\left(\eta^{\sigma-1}e^{-\frac{\pi t}{2}-\frac{t}{16}}\frac{\eta}{\sqrt{t}}\left(1 + \sum_{n=1}^{N-1}\left(\frac{5et}{16n}\right)^{\frac{n}{3}}\right)\right). \end{aligned}$$

Now $(t/n)^{\frac{n}{3}}$ increases steadily up to $n = t/e$, and so if $n < At$, where $A < 1/e$, it is

$$O(e^{\frac{1}{3}tA\log 1/A}).$$

Hence if $N < At$, with A sufficiently small, the total error is

$$O(e^{-\frac{\pi t}{2}-At}).$$

This proves (4.38).

Finally, we analyze the sum

$$(i\eta)^{s-1}\sum_{n=0}^{N-1} a_n \int_{C_2'} \frac{e^{\frac{t}{\eta}(w-i\eta)+\frac{it}{2\eta^2}(w-i\eta)^2-mw}}{e^w-1}(w-i\eta)^n dw. \tag{4.42}$$

The integral in (4.42) may be expressed as

$$-\int_L e^{\frac{t}{\eta}(w+2[\frac{\eta}{2\pi}]\pi i-i\eta)+\frac{it}{2\eta^2}(w+2[\frac{\eta}{2\pi}]\pi i-i\eta)^2-[\frac{t}{\eta}]w}\frac{(w+2[\frac{\eta}{2\pi}]\pi i-i\eta)^n}{e^w-1}dw,$$

where L is a line in the direction $\arg w = \pi/4$, passing between 0 and $2\pi i$. This is $n!$ times the coefficient of ξ^n in the following expression:

$$\begin{aligned} &-\int_L e^{\frac{t}{\eta}(w+2[\frac{\eta}{2\pi}]\pi i-i\eta)+\frac{it}{2\eta^2}(w+2[\frac{\eta}{2\pi}]\pi i-i\eta)^2-[\frac{t}{\eta}]w+\xi(w+2[\frac{\eta}{2\pi}]\pi i-i\eta)}\frac{dw}{e^w-1} \\ &= -e^{\frac{t}{\eta}2[\frac{\eta}{2\pi}]\pi i-it+\frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi i-i\eta)^2+\xi(2[\frac{\eta}{2\pi}]\pi i-i\eta)}\int_L e^{\frac{it}{2\eta^2}w^2+(\frac{2t}{\eta}-\frac{2\pi t}{\eta^2}[\frac{\eta}{2\pi}]-[\frac{t}{\eta}]+\xi)w}\frac{dw}{e^w-1} \\ &= e^{\frac{t}{\eta}2[\frac{\eta}{2\pi}]\pi i-it+\frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi i-i\eta)^2}2\pi i\Phi\left(-\frac{2\pi t}{\eta^2}, \frac{2t}{\eta}-\frac{2\pi t}{\eta^2}\left[\frac{\eta}{2\pi}\right]-\left[\frac{t}{\eta}\right]+\xi-\frac{1}{2}\right)e^{\xi(2[\frac{\eta}{2\pi}]\pi i-i\eta)}, \end{aligned} \tag{4.43}$$

where the function $\Phi(\tau, u)$ is defined by (4.28), i.e.

$$\Phi(\tau,u) = \int_{0\nwarrow 1}\frac{e^{\pi i\tau x^2+2\pi iux}}{e^{\pi ix}-e^{-\pi ix}}dx = \frac{1}{2\pi i}\int_{-L}\frac{e^{-\frac{i\tau}{4\pi}w^2+(u+\frac{1}{2})w}}{e^w-1}dw, \qquad \tau<0, \quad u\in\mathbb{C}.$$

We rewrite the expression on the rhs of (4.43) as

$$\begin{aligned} e^{\frac{t}{\eta}2[\frac{\eta}{2\pi}]\pi i-it+\frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi i-i\eta)^2}2\pi i\sum_{l=0}^{\infty}\partial_2^l\Phi\left(-\frac{2\pi t}{\eta^2}, \frac{2t}{\eta}-\frac{2\pi t}{\eta^2}\left[\frac{\eta}{2\pi}\right]-\left[\frac{t}{\eta}\right]-\frac{1}{2}\right)\frac{\xi^l}{l!} \\ \times\sum_{k=0}^{\infty}\frac{\xi^k(2[\frac{\eta}{2\pi}]\pi i-i\eta)^k}{k!}. \end{aligned}$$

Hence the expression in (4.42) is given by

$$2\pi i(i\eta)^{s-1}e^{\frac{t}{\eta}2[\frac{\eta}{2\pi}]\pi i-it+\frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi i-i\eta)^2}S_N(s,\eta) \tag{4.44}$$

where $S_N(s,\eta)$ is defined in (4.26). Equations (4.36), (4.38), and (4.44) show that I_2 satisfies

$$I_2 = 2\pi i(i\eta)^{s-1} e^{\frac{t}{\eta}2[\frac{\eta}{2\pi}]\pi i - it + \frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi i - i\eta)^2} S_N(s,\eta) + O\left(e^{-\frac{\pi t}{2}} \left(\frac{3N}{t}\right)^{\frac{N}{6}} \frac{\eta^\sigma}{\sqrt{t}}\right) + O(e^{-\frac{\pi t}{2} - At}).$$

Equation (4.25) follows by substituting this expression into (4.3).

In order to prove (4.29) we suppose that $u \in \mathbb{C}$ and $\tau < 0$. We claim that Φ satisfies the two recursion relations

$$\Phi(\tau, u) = \Phi(\tau, u+1) - \frac{e^{\frac{3\pi i}{4}}}{\sqrt{|\tau|}} e^{-\frac{\pi i}{\tau}(u+\frac{1}{2})^2} \tag{4.45}$$

and

$$\Phi(\tau, u) = 1 - e^{\pi i \tau - 2\pi i u} \Phi(\tau, u - \tau). \tag{4.46}$$

Repetitive use of these two equations yields the following identities:

$$\Phi(\tau, u) = \Phi(\tau, u+N) - \frac{e^{\frac{3\pi i}{4}}}{\sqrt{|\tau|}} \sum_{n=0}^{N-1} e^{-\frac{\pi i}{\tau}(u+n+\frac{1}{2})^2}, \qquad N \geq 0, \tag{4.47}$$

and

$$\Phi(\tau, u) = \sum_{n=0}^{N-1} (-1)^n e^{\pi i n^2 \tau - 2\pi i n u} + (-1)^N e^{\pi i N^2 \tau - 2\pi i N u} \Phi(\tau, u - N\tau), \qquad N \geq 0. \tag{4.48}$$

If $\tau = -p/q$, we apply (4.47) with $N = p$ and (4.48) with $N = q$ to find

$$\Phi(\tau, u) = \Phi(\tau, u+p) - \frac{e^{\frac{3\pi i}{4}}}{\sqrt{|\tau|}} \sum_{n=0}^{p-1} e^{-\frac{\pi i}{\tau}(u+n+\frac{1}{2})^2} \tag{4.49}$$

and

$$\Phi(\tau, u) = \sum_{n=0}^{q-1} (-1)^n e^{\pi i n^2 \tau - 2\pi i n u} + (-1)^q e^{\pi i q^2 \tau - 2\pi i q u} \Phi(\tau, u+p). \tag{4.50}$$

Eliminating $\Phi(\tau, u+p)$ from these two equations, we find (4.29).

It remains to prove (4.45) and (4.46). In order to prove (4.45), we note that

$$\begin{aligned} \Phi(\tau, u+1) - \Phi(\tau, u) &= \int_{0 \nwarrow 1} e^{\pi i \tau x^2} \frac{e^{2\pi i (u+1)x} - e^{2\pi i u x}}{e^{\pi i x} - e^{-\pi i x}} dx \\ &= \int_{0 \nwarrow 1} e^{\pi i \tau x^2 + 2\pi i (u+\frac{1}{2})x} dx \end{aligned}$$

$$\begin{aligned}
&= e^{-\frac{\pi i}{\tau}(u+\frac{1}{2})^2}\int_{0\nwarrow 1} e^{\pi i\tau(x+\frac{u+\frac{1}{2}}{\tau})^2}dx \\
&= e^{-\frac{\pi i}{\tau}(u+\frac{1}{2})^2}\int_{0\nwarrow 1} e^{\pi i\tau x^2}dx.
\end{aligned}$$

The identities

$$\int_{0\nwarrow 1} e^{\pi i\tau x^2}dx = \int_{0\nwarrow 1} e^{-\pi i y^2}\frac{dy}{\sqrt{|\tau|}} = \frac{e^{\frac{3\pi i}{4}}}{\sqrt{|\tau|}}$$

imply equation (4.45).

The identity (4.46) is established as follows:

$$\begin{aligned}
\Phi(\tau,u) &= \int_{0\nwarrow 1}\frac{e^{\pi i\tau x^2+2\pi iux}}{e^{\pi ix}-e^{-\pi ix}}dx \\
&= 1+\int_{-1\nwarrow 0}\frac{e^{\pi i\tau x^2+2\pi iux}}{e^{\pi ix}-e^{-\pi ix}}dx \\
&= 1+\int_{0\nwarrow 1}\frac{e^{\pi i\tau(x-1)^2+2\pi iu(x-1)}}{e^{\pi i(x-1)}-e^{-\pi i(x-1)}}dx \\
&= 1-e^{\pi i\tau-2\pi iu}\int_{0\nwarrow 1}\frac{e^{\pi i\tau x^2+2\pi i(u-\tau)x}}{e^{\pi ix}-e^{-\pi ix}}dx \\
&= 1-e^{\pi i\tau-2\pi iu}\Phi(\tau,u-\tau),
\end{aligned}$$

where the second equality follows from the fact that the residue of the integrand at $x=0$ is $\frac{1}{2\pi i}$. **QED**

Theorem 4.4 is valid for $\epsilon\sqrt{t}<\eta<t$. As a corollary, we can find an analogous result valid for $2\pi<\eta<\frac{\sqrt{t}}{\epsilon}$.

Corollary 4.6 (The asymptotics to all orders for the case $2\pi<\eta<\frac{\sqrt{t}}{\epsilon}$) *For every $\epsilon>0$, there exists a constant $A>0$ such that*

$$\begin{aligned}
\zeta(s) = &\sum_{n=1}^{[\frac{t}{\eta}]}\frac{1}{n^s}+\chi(s)\sum_{n=1}^{[\frac{\eta}{2\pi}]}\frac{1}{n^{1-s}} \qquad (4.51)\\
&+\chi(s)e^{i\pi(1-s)}\Gamma(s)\Bigg\{e^{\frac{i\pi s}{2}}\left(\frac{\eta}{2\pi t}\right)^s e^{-i\eta[\frac{t}{\eta}]+it+\frac{i\eta^2}{2t}([\frac{t}{\eta}]-\frac{t}{\eta})^2}\overline{S_N\left(1-\bar{s},\frac{2\pi t}{\eta}\right)} \\
&\qquad+O\left(e^{-\frac{\pi t}{2}}\left(\frac{3N}{t}\right)^{\frac{N}{6}}\frac{(\frac{2\pi t}{\eta})^{1-\sigma}}{\sqrt{t}}\right)\Bigg\},
\end{aligned}$$

$$2\pi<\eta<\frac{\sqrt{t}}{\epsilon},\quad 0\le\sigma\le 1,\quad 1\le N<At,\quad t\to\infty,$$

where the error term is uniform for all η,σ,N in the above ranges and S_N is given by (4.26).

Proof. We replace σ by $1-\sigma$ in (4.25) and take the complex conjugate of both sides. We then multiply the resulting equation by $\chi(s)$ and use the identities (4.23). This yields the following equation

$$\begin{aligned}\zeta(s) &= \chi(s)\sum_{n=1}^{[\frac{t}{\eta}]}\frac{1}{n^{1-s}}+\sum_{n=1}^{[\frac{\eta}{2\pi}]}\frac{1}{n^{s}} \qquad (4.52)\\ &+\chi(s)e^{i\pi(1-s)}\Gamma(s)\Bigg\{(e^{\frac{i\pi s}{2}}\eta^{-s}e^{-\frac{2t}{\eta}[\frac{\eta}{2\pi}]\pi i+it+\frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi-\eta)^2}\overline{S_N(1-\bar{s},\eta)}\\ &\qquad +O\Bigg(e^{-\frac{\pi t}{2}}\left(\frac{3N}{t}\right)^{\frac{N}{6}}\frac{\eta^{1-\sigma}}{\sqrt{t}}\Bigg)\Bigg\},\\ &\epsilon\sqrt{t}<\eta<t,\quad 0\le\sigma\le 1,\quad 1\le N<At,\quad t\to\infty.\end{aligned}$$

Replacing η by $\frac{2\pi t}{\eta}$, we find (4.51). **QED**

Remark 4.7 The results of theorem 4.4 are convenient for studying the higher-order asymptotics of $\zeta(s)$ in the case when $\eta = \text{constant}\times\sqrt{t}$. However, if the order of η is strictly smaller or larger than the order of $\sqrt{t}$, the results of theorem 4.4 are less convenient, because in this case the asymptotics of $\zeta(s)$ involves the asymptotics of the sums in (4.29) as p and/or q tend to infinity. In this case, the alternative representation of the asymptotics of theorem 4.1, which avoids the appearance of the above sums, is usually more convenient.

5 Consequences of the Asymptotic Formulae

Using theorems 3.1 and 3.2 we can compute several interesting sums.

Theorem 5.1 *Define the polylogarithm* $\mathrm{Li}_m(z)$ *by (3.3). The following relation holds:*

$$\begin{aligned}\sum_{n=[\frac{\eta_1}{2\pi}]+1}^{[\frac{\eta_2}{2\pi}]} n^{-s} &= \frac{1}{1-s}\left[\left(\frac{\eta_2}{2\pi}\right)^{1-s}-\left(\frac{\eta_1}{2\pi}\right)^{1-s}\right] \qquad (5.1)\\ &+\frac{e^{-\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}}\sum_{n=1}^{\infty}\sum_{j=0}^{N-1}e^{-nz-it\ln z}\left(\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\right)^{j}\frac{z^{-\sigma}}{n+\frac{it}{z}}\Bigg|_{z=i\eta_2}^{i\eta_1}\\ &+\frac{e^{\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}}\sum_{n=1}^{\infty}\sum_{j=0}^{N-1}e^{-nz-it\ln z}\left(\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\right)^{j}\frac{z^{-\sigma}}{n+\frac{it}{z}}\Bigg|_{z=-i\eta_2}^{-i\eta_1}\\ &+O\Bigg((2N+1)!!N\Big(\frac{1+\epsilon}{\epsilon}\Big)^{2(N+1)}\eta_1^{-\sigma-N}\Bigg),\\ &(1+\epsilon)t<\eta_1<\eta_2<\infty,\quad \epsilon>0,\quad 0\le\sigma\le1,\quad N\ge2,\quad t\to\infty,\end{aligned}$$

where the error term is uniform for all $\eta_1, \eta_2, \epsilon, \sigma, N$ *in the above ranges.*

For $N = 3$ *equation (5.1) simplifies to*

$$\begin{aligned}
\sum_{n=[\frac{\eta_1}{2\pi}]+1}^{[\frac{\eta_2}{2\pi}]} n^{-s} = {} & \frac{1}{1-s}\left[\left(\frac{\eta_2}{2\pi}\right)^{1-s} - \left(\frac{\eta_1}{2\pi}\right)^{1-s}\right] \qquad (5.2)\\
& + \frac{2i\eta_1^{-s}}{(2\pi)^{1-s}}\Big\{-i\arg(1-e^{i\eta_1}) + \frac{t-i\sigma}{\eta_1}\operatorname{Re}\operatorname{Li}_2(e^{i\eta_1})\\
& \qquad + \frac{1}{\eta_1^2}\big[it^2 - 3i\sigma t - (\sigma-1)t - i\sigma(\sigma+1)\big]\operatorname{Im}\operatorname{Li}_3(e^{i\eta_1})\Big\}\\
& - \frac{2i\eta_2^{-s}}{(2\pi)^{1-s}}\Big\{-i\arg(1-e^{i\eta_2}) + \frac{t-i\sigma}{\eta_2}\operatorname{Re}\operatorname{Li}_2(e^{i\eta_2})\\
& \qquad + \frac{1}{\eta_2^2}\big[it^2 - 3i\sigma t - (\sigma-1)t - i\sigma(\sigma+1)\big]\operatorname{Im}\operatorname{Li}_3(e^{i\eta_2})\Big\}\\
& + O\left(\frac{1}{\eta_1^{3+\sigma}}\left(t^3 + \left(\frac{1+\epsilon}{\epsilon}\right)^8\right)\right),\\
& \qquad (1+\epsilon)t < \eta_1 < \eta_2 < \infty, \quad \epsilon > 0, \quad 0 \le \sigma \le 1, \quad t \to \infty,
\end{aligned}$$

where the error term is uniform for all $\eta_1, \eta_2, \epsilon, \sigma$ *in the above ranges.*

Proof. Replacing in equation (3.1) η with η_1 and subtracting the resulting equation from the equation obtained from (3.1) by replacing η with η_2, we find (5.1). Equation (5.2) follows in a similar way from (3.2). **QED**

Theorem 5.2 *Define the polylogarithm* $\operatorname{Li}_m(z)$ *by (3.3). The following relation holds:*

$$\begin{aligned}
\sum_{n=[\frac{t}{2\pi}]+1}^{[\frac{\eta}{2\pi}]} n^{-s} = {} & \frac{1}{1-s}\left[\left(\frac{\eta}{2\pi}\right)^{1-s} - \left(\frac{t}{2\pi}\right)^{1-s}\right] \qquad (5.3)\\
& + \frac{e^{-\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}}\sum_{n=1}^{\infty}\sum_{j=0}^{N-1} e^{-nz-it\ln z}\left(\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\right)^j \frac{z^{-\sigma}}{n+\frac{it}{z}}\Bigg|_{z=i\eta}^{it}\\
& - \frac{e^{\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}}\sum_{n=1}^{\infty}\sum_{j=0}^{N-1} e^{-nz-it\ln z}\left(\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\right)^j \frac{z^{-\sigma}}{n+\frac{it}{z}}\Bigg|_{z=-i\eta}\\
& + \frac{e^{\frac{i\pi(1-s)}{2}}}{(2\pi)^{1-s}}\sum_{n=2}^{\infty}\sum_{j=0}^{N-1} e^{-nz-it\ln z}\left(\frac{1}{n+\frac{it}{z}}\frac{d}{dz}\right)^j \frac{z^{-\sigma}}{n+\frac{it}{z}}\Bigg|_{z=-it}\\
& + \left(\frac{t}{2\pi}\right)^{1-s} e^{it}\sum_{k=0}^{2N}\frac{\overline{c_k(1-\sigma)\Gamma(\frac{k+1}{2})}}{t^{\frac{k+1}{2}}}
\end{aligned}$$

$$+\,O\biggl(\frac{(2N+1)!!Ne^{2(N+1)}}{t^{\sigma+N}}+\frac{(2N+1)!!N(\frac{1+\epsilon}{\epsilon})^{2(N+1)}}{\eta^{\sigma+N}}\biggr),$$

$$(1+\epsilon)t<\eta<\infty,\quad \epsilon>0,\quad 0\le\sigma\le 1,\quad N\ge 2,\quad t\to\infty,$$

where the error term is uniform for all $\eta, \epsilon, \sigma, N$ *in the above ranges and the coefficients* $c_k(\sigma)$ *are defined in (3.30).*

For $N=3$ *equation (5.3) simplifies to*

$$\sum_{n=[\frac{t}{2\pi}]+1}^{[\frac{\eta}{2\pi}]} n^{-s}=\frac{1}{1-s}\biggl[\Bigl(\frac{\eta}{2\pi}\Bigr)^{1-s}-\Bigl(\frac{t}{2\pi}\Bigr)^{1-s}\biggr]+\Bigl\{\cdots\Bigr\} \tag{5.4}$$

$$+\,O\biggl(\frac{1}{\eta^{3+\sigma}}\Bigl(t^3+\Bigl(\frac{1+\epsilon}{\epsilon}\Bigr)^8\Bigr)+\frac{1}{t^{3+\sigma}}\biggr),$$

$$(1+\epsilon)t<\eta<\infty,\quad \epsilon>0,\quad 0\le\sigma\le 1,\quad t\to\infty,$$

where the error term is uniform for all η, ϵ, σ *in the above ranges and* $\{\cdots\}$ *denotes an explicit expression involving* σ*,* t*, and the polylogarithms* $\mathrm{Li}_m(e^{it})$*,* $m=1,\ldots,5$*.*

Proof. Equation (5.3) follows by subtracting equation (3.20) from equation (3.1). Similarly, equation (5.4) follows by subtracting equation (3.22) from equation (3.2).

QED

The following equation is given on page 78 of Titchmarsh [9]:

$$\sum_{x<n\le N}\frac{1}{n^s}\sim\chi(s)\sum_{\frac{t}{2\pi N}<n\le\frac{t}{2\pi x}}\frac{1}{n^{1-s}}. \tag{5.5}$$

Theorem 4.4 yields a precise version of the relation (5.5) with an explicit error estimate.

Theorem 5.3 *For every* $\epsilon>0$*, there exists a constant* $A>0$ *such that*

$$\sum_{n=[\frac{t}{\eta_2}]+1}^{[\frac{t}{\eta_1}]}\frac{1}{n^s}=\chi(s)\sum_{n=[\frac{\eta_1}{2\pi}]+1}^{[\frac{\eta_2}{2\pi}]}\frac{1}{n^{1-s}} \tag{5.6}$$

$$+\,e^{-i\pi s}\Gamma(1-s)\biggl\{e^{\frac{i\pi(s-1)}{2}}\eta^{s-1}e^{\frac{2t}{\eta}[\frac{\eta}{2\pi}]\pi i-it-\frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi-\eta)^2}S_N(s,\eta)$$

$$+\,O\biggl(e^{-\frac{\pi t}{2}}\Bigl(\frac{3N}{t}\Bigr)^{\frac{N}{6}}\frac{\eta^\sigma}{\sqrt{t}}\biggr)\biggr\}\bigg|_{\eta=\eta_1}^{\eta_2},$$

$$\epsilon\sqrt{t}<\eta_1<\eta_2<t,\quad 0\le\sigma\le 1,\quad 1\le N<At,\quad t\to\infty,$$

where the error term is uniform for all $\eta_1, \eta_2, \sigma, N$ *in the above ranges and the function* $S_N(s,\eta)$ *is defined by (4.26).*

For $N = 3$ equation (5.6) simplifies to

$$\begin{aligned}
\sum_{n=[\frac{t}{\eta_2}]+1}^{[\frac{t}{\eta_1}]} \frac{1}{n^s} = \chi(s) \sum_{n=[\frac{\eta_1}{2\pi}]+1}^{[\frac{\eta_2}{2\pi}]} \frac{1}{n^{1-s}} \qquad\qquad (5.7)\\
+ e^{-i\pi s}\Gamma(1-s)\Bigg\{ e^{\frac{i\pi(s-1)}{2}} \eta^{s-1} e^{\frac{t}{\eta}2[\frac{\eta}{2\pi}]\pi i - it - \frac{it}{2\eta^2}(2[\frac{\eta}{2\pi}]\pi-\eta)^2} \\
\times \Bigg[\Phi + \frac{\sigma-1}{i\eta}\left(\partial_2\Phi + \left(2\left[\frac{\eta}{2\pi}\right]\pi i - i\eta \right)\Phi \right) \\
- \frac{(\sigma-2)(\sigma-1)}{2\eta^2}\left(\partial_2^2\Phi + 2\left(2\left[\frac{\eta}{2\pi}\right]\pi i - i\eta \right)\partial_2\Phi \right. \\
\left. + \left(2\left[\frac{\eta}{2\pi}\right]\pi i - i\eta \right)^2 \Phi \right) \Bigg] + O\left(e^{-\frac{\pi t}{2}} \frac{\eta^\sigma}{t} \right) \Bigg\} \Bigg|_{\eta=\eta_1}^{\eta_2}, \\
\epsilon\sqrt{t} < \eta_1 < \eta_2 < t, \quad 0 \le \sigma \le 1, \quad t \to \infty,
\end{aligned}$$

where the error term is uniform for all η_1, η_2, σ in the above ranges, Φ is defined by (4.28), and Φ and its partial derivatives are evaluated at the point (1.8).

Proof. Replacing in equation (4.25) η with η_1 and subtracting the resulting equation from the equation obtained from (4.25) by replacing η with η_2, we find (5.6). Equation (5.7) follows in a similar way from (4.30). **QED**

Remark 5.4 The relation (5.5) is a particular case of (5.6) (let $x = t/\eta_2$ and $N = t/\eta_1$).

6 Several Representations for the Basic Sum

The goal of this section is to present integral representations of the basic sum $\sum_a^b n^{s-1}$ for certain values of a and b.

Let the contour C_t^η, $t < \eta$, denote the semicircle from it to $i\eta$ with $\operatorname{Re} z \ge 0$. Splitting the contour of the second integral in the rhs of equation (1.2) into the contour C_η^t plus the ray from it to $\infty \exp(i\phi_2)$, we find

$$\begin{aligned}
\zeta(1-s) = \sum_{n=1}^{[\frac{\eta}{2\pi}]} n^{s-1} - \frac{\eta^s}{s(2\pi)^s} - \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s}\int_{C_t^\eta} \frac{z^{s-1}}{e^z-1}dz \\
+ G_L(t,\sigma;\eta) + G_U(t,\sigma;t), \qquad 0 \le \sigma \le 1, \quad 0 < t < \eta, \qquad (6.1)
\end{aligned}$$

where G_L and G_U are defined in (3.5) and (3.6) respectively. The term $G_U(t,\sigma;t)$ was computed to all orders as $t \to \infty$ in theorem 3.2, see equations (3.26) and (3.33). The term $G_L(t,\sigma;\eta)$ with $t<\eta$ was computed to all orders as $t\to\infty$ in (3.13). Thus, by comparing equation (6.1) with the representation obtained in theorem 3.2, it follows that we can express the fundamental sum $\sum_{n=[t/2\pi]+1}^{[\eta/2\pi]} n^{s-1}$ in terms of the integral appearing in the rhs of (6.1), where the error is computed to all orders. For brevity of presentation, we state this result only to leading order.

Lemma 6.1 (An integral representation of the basic sum) *Let C_t^η, $t<\eta$, denote the semicircle from $z=it$ to $z=i\eta$ with $\operatorname{Re} z\geq 0$. For every $\epsilon>0$,*

$$\begin{aligned}\sum_{n=[\frac{t}{2\pi}]+1}^{[\frac{\eta}{2\pi}]} n^{s-1} = {} & \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s}\int_{C_t^\eta}\frac{z^{s-1}}{e^z-1}dz+\frac{\eta^s}{s(2\pi)^s}+\frac{i\eta^{s-1}}{(2\pi)^s}\log(1-e^{i\eta}) \\ & -\frac{it^{s-1}e^{-it}}{(2\pi)^s}\log(1-e^{it})+O\left(\frac{1}{t^{2-\sigma}}+\frac{t}{\eta^{2-\sigma}}\right), \\ & (1+\epsilon)t<\eta<\infty, \quad 0\leq\sigma\leq 1, \quad t\to\infty,\end{aligned} \tag{6.2}$$

where the error term is uniform for all η,σ in the above ranges.

Proof. Letting $\sigma\to 1-\sigma$ in (3.22) and taking the complex conjugate of the resulting equation, we find

$$\begin{aligned}\zeta(1-s) = {} & \sum_{n=1}^{[\frac{t}{2\pi}]} n^{s-1}-\frac{1}{s}\left(\frac{t}{2\pi}\right)^s+\frac{it^{s-1}e^{-it}}{(2\pi)^s}(-e^{it}+2i\operatorname{Im}\operatorname{Li}_1(e^{it})) \\ & +\frac{t^{s-1}e^{-it}}{(2\pi)^s}\left(\frac{1-i}{2}\sqrt{\pi t}+\frac{i}{3}-i\sigma-\frac{1+i}{24}\frac{\sqrt{\pi}\,(6\sigma^2-6\sigma+1)}{\sqrt{t}}\right)+O(t^{\sigma-2}).\end{aligned}$$

The lemma follows by subtracting this equation from (6.1) and using the following equations which follow from (3.13), (3.33), and (3.26) respectively:

$$\begin{aligned}G_L(t,\sigma;\eta) &= -\frac{i\eta^{s-1}}{(2\pi)^s}\log(1-e^{i\eta})+O\left(\frac{t}{\eta^{2-\sigma}}\right), \quad (1+\epsilon)t<\eta, \text{ (not uniformly in } \epsilon), \\ G_U^{(1)}(t,\sigma;t) &= \frac{t^{s-1}e^{-it}}{(2\pi)^s}\left(\frac{1-i}{2}\sqrt{\pi t}+\frac{i}{3}-i\sigma-\frac{1+i}{24}\frac{\sqrt{\pi}\,(6\sigma^2-6\sigma+1)}{\sqrt{t}}\right)+O(t^{\sigma-2}), \\ G_U^{(2)}(t,\sigma;t) &= \frac{it^{s-1}e^{-it}}{(2\pi)^s}\log(1-e^{-it})+O(t^{\sigma-2}).\end{aligned}$$

QED

It is possible to derive an alternative integral representation of the related sum $\sum_{[\eta/2\pi]+1}^{[t/2\pi]} n^{s-1}$.

Lemma 6.2 (An alternative integral representation of the basic sum) *For every $\delta > 0$,*

$$\sum_{n=[\frac{\eta}{2\pi}]+1}^{[\frac{t}{2\pi}]} n^{s-1} = \frac{2}{(2\pi)^s} \fint_{\eta}^{t} \frac{\rho^{s-1}}{e^{i\rho}-1} d\rho + O(t^{\sigma-1}), \tag{6.3}$$

$$0 < \eta < t, \quad \operatorname{dist}(\eta, 2\pi\mathbb{Z}) > \delta, \quad \operatorname{dist}(t, 2\pi\mathbb{Z}) > \delta, \quad 0 \le \sigma \le 1, \quad t \to \infty,$$

where the error term is uniform for all η, σ in the above ranges and the contour in the integral denotes the principal value integral with respect to the points

$$\left\{ 2\pi n \,\middle|\, n \in \mathbb{Z}^+, \; \left[\frac{\eta}{2\pi}\right] + 1 \le n \le \left[\frac{t}{2\pi}\right] \right\}.$$

Proof. Let $\hat{C}_t^\eta$ denote the semicircle from it to $i\eta$ with $\operatorname{Re} z \le 0$, defined in equation (2.12c) with η and α replaced by t and η respectively, see figure 6. Let $R_\eta^t(s)$ and $L_t^\eta(s)$ denote the following:

$$R_\eta^t(s) = \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s} \int_{C_\eta^t} \frac{z^{s-1}}{e^z - 1} dz, \qquad s \in \mathbb{C} \tag{6.4}$$

and

$$L_t^\eta(s) = \frac{e^{-\frac{i\pi s}{2}}}{(2\pi)^s} \int_{\hat{C}_t^\eta} \frac{z^{s-1}}{e^z - 1} dz, \qquad s \in \mathbb{C}. \tag{6.5}$$

Cauchy's theorem applied in the interior of the disk whose boundary is $C_\eta^t \cup \hat{C}_t^\eta$, yields

$$L_t^\eta(s) + R_\eta^t(s) = \sum_{n=[\frac{\eta}{2\pi}]+1}^{[\frac{t}{2\pi}]} n^{s-1}, \qquad s \in \mathbb{C}. \tag{6.6}$$

The Plemelj formulas imply

$$R_\eta^t - L_t^\eta = \frac{2}{(2\pi)^s} \fint_{\eta}^{t} \frac{\rho^{s-1}}{e^{i\rho} - 1} d\rho. \tag{6.7}$$

Using (6.7) to replace R_η^t in (6.6), we find

$$\sum_{n=[\frac{\eta}{2\pi}]+1}^{[\frac{t}{2\pi}]} n^{s-1} = \frac{2}{(2\pi)^s} \fint_{\eta}^{t} \frac{\rho^{s-1}}{e^{i\rho} - 1} d\rho + 2L_t^\eta. \tag{6.8}$$

Replacing the contour $\hat{C}_t^\eta$ in (6.5) by the union of following three segments:

$$\{ite^{i\theta} \mid 0 \le \theta \le \pi/4\}, \quad \{i\rho e^{\frac{\pi i}{4}} \mid \eta \le \rho \le t\}, \quad \{i\eta e^{i\theta} \mid 0 \le \theta \le \pi/4\},$$



Figure 6 *The contours* C_η^t *and* $\hat{C}_t^\eta$.

we obtain

$$\begin{aligned}L_t^\eta &= \frac{1}{(2\pi)^s}\left[\int_0^{\frac{\pi}{4}} \frac{e^{i\theta(s-1)}t^{s-1}}{e^{ite^{i\theta}}-1} ite^{i\theta}d\theta - \int_\eta^t \frac{e^{\frac{\pi i}{4}(s-1)}\rho^{s-1}}{e^{i\rho e^{\frac{\pi i}{4}}}-1} e^{\frac{\pi i}{4}}d\rho - \int_0^{\frac{\pi}{4}} \frac{e^{i\theta(s-1)}\eta^{s-1}}{e^{i\eta e^{i\theta}}-1} i\eta e^{i\theta}d\theta\right] \\ &=: \frac{1}{(2\pi)^s}[J_1 + J_2 + J_3].\end{aligned}$$

The assumption $\operatorname{dist}(t, 2\pi\mathbb{Z}) > \delta$ implies that

$$|J_1| \leq \int_0^{\frac{\pi}{4}} \frac{e^{-\theta t}t^{\sigma-1}}{A} td\theta = \frac{t^{\sigma-1}}{A}(1 - e^{-\frac{\pi t}{4}}) = O(t^{\sigma-1}).$$

Similar computations show that J_2 is exponentially small and, in view of the assumption $\operatorname{dist}(\eta, 2\pi\mathbb{Z}) > \delta$, that J_3 is $O(\eta^\sigma t^{-1})$. Thus,

$$L_t^\eta = O(t^{\sigma-1}), \tag{6.9}$$

and the lemma follows from (6.8). **QED**

Remark 6.3 Lemma 6.2 implies that the leading behavior of $\zeta(s)$ is characterized by the following integral:

$$I(\eta, t, \sigma) = \fint_\eta^t \frac{\rho^{s-1}}{e^{i\rho}-1} d\rho.$$

Remark 6.4 The estimate (6.9) is a consequence of the fact that the integral appearing in the definition of $L_t^\eta(s)$ does *not* possess any stationary points. Indeed, using

$$\frac{1}{e^z - 1} = -\sum_{m=0}^{\infty} e^{mz}, \qquad \operatorname{Re} z < 0,$$

it follows that

$$e^{-\frac{i\pi s}{2}}\int_{\hat{C}_t^\eta}\frac{z^{s-1}}{e^z-1}dz=\sum_{m=0}^{\infty}\int_\eta^t e^{im\rho+it\ln\rho}\rho^{\sigma-1}d\rho.$$

Candidates for stationary points occur at $\rho^*=-t/m$ and the inequality $\eta\le\rho^*\le t$ implies the non-existence of any stationary points.

The next lemma gives an alternative representation for the fundamental integral $R_\eta^t(s)$ defined in (6.4).

Lemma 6.5 *Let C_η^t denote the semicircle from $i\eta$ to it with* $\operatorname{Re} z\ge 0$ *defined by equation (1.12). For every $\delta>0$,*

$$R_\eta^t(s)=\frac{e^{-i\epsilon s}}{(2\pi)^s}\int_\eta^t\frac{u^{s-1}du}{e^{iue^{-i\epsilon}}-1}+O\left(\frac{e^{\epsilon t}-1}{t^{1-\sigma}}\right),\tag{6.10}$$

$$0<\eta<t,\quad \operatorname{dist}(\eta,2\pi\mathbb{Z})>\delta,\quad \operatorname{dist}(t,2\pi\mathbb{Z})>\delta,\quad 0\le\sigma\le1,\quad 0<\epsilon<1,\quad t\to\infty,$$

where the error term is uniform for all η,σ,ϵ in the above ranges.

Proof. Let $\epsilon\in(0,1)$. We deform the contour C_η^t on the lhs of (6.10) so that it consists of the following three pieces:

$$\{i\eta e^{-i\theta}\mid 0\le\theta\le\epsilon\},\quad\{iue^{-i\epsilon}\mid\eta\le u\le t\},\quad\{ite^{-i\theta}\mid 0\le\theta\le\epsilon\}.$$

This yields

$$e^{-\frac{i\pi s}{2}}\int_{C_\eta^t}\frac{z^{s-1}dz}{e^z-1}=e^{-i\epsilon s}\int_\eta^t\frac{u^{s-1}du}{e^{iue^{-i\epsilon}}-1}-i\eta^s\int_0^\epsilon\frac{e^{-i\theta s}d\theta}{e^{i\eta e^{-i\theta}}-1}+it^s\int_0^\epsilon\frac{e^{-i\theta s}d\theta}{e^{ite^{-i\theta}}-1}.$$

The assumption that $\operatorname{dist}(\eta,2\pi\mathbb{Z})>\delta$ implies that there exists an A such that $|e^{i\eta e^{-i\theta}}-1|\ge A$ for all $\theta\in[0,\epsilon]$, $\epsilon\in(0,1)$, and $\eta>0$. Thus,

$$\left|\frac{e^{-i\theta s}}{e^{i\eta e^{-i\theta}}-1}\right|\le\frac{e^{\theta t}}{A},\qquad\theta\in[0,\epsilon],$$

and so

$$\left|\int_0^\epsilon\frac{e^{-i\theta s}}{e^{i\eta e^{-i\theta}}-1}d\theta\right|\le\frac{e^{\epsilon t}-1}{At}.$$

Similarly, because of the assumption $\operatorname{dist}(t,2\pi\mathbb{Z})>\delta$,

$$\left|\int_0^\epsilon\frac{e^{-i\theta s}}{e^{ite^{-i\theta}}-1}d\theta\right|\le\frac{e^{\epsilon t}-1}{At}.$$

This proves the lemma. **QED**

Lemma 6.5 implies that

$$R_\eta^t(s) = \frac{1}{(2\pi)^s} \lim_{\substack{\epsilon\to 0\\ \epsilon t\to 0}} \int_\eta^t \frac{u^{s-1}du}{e^{iue^{-i\epsilon}}-1}, \tag{6.11}$$

which in view of (6.6) and (6.9) implies the following alternative representation for the basic sum:

$$\frac{1}{(2\pi)^s} \lim_{\substack{\epsilon\to 0\\ \epsilon t\to 0}} \int_\eta^t \frac{u^{s-1}du}{e^{iue^{-i\epsilon}}-1} = \sum_{n=[\frac{\eta}{2\pi}]+1}^{[\frac{t}{2\pi}]} n^{s-1} + O(t^{\sigma-1}).$$

In fact, in the remaining part of this section we will present arguments which suggest[2] that on the critical line the assumption $\epsilon t \to 0$ in (6.11) can be relaxed to $\epsilon^2 t \to 0$, so that the following equation is valid:

$$R_\eta^t(s) = \frac{1}{(2\pi)^s} \lim_{\substack{\epsilon\to 0\\ \epsilon^2 t\to 0}} \int_\eta^t \frac{u^{s-1}du}{e^{iue^{-i\epsilon}}-1}, \tag{6.12}$$

where it is assumed that $\mathrm{dist}(\eta, 2\pi\mathbb{Z}) > \delta$, $\eta = O(1)$, $\mathrm{dist}(t, 2\pi\mathbb{Z}) > \delta$, and $\sigma = 1/2$.

Indeed, consider the integral

$$(2\pi)^s R_\eta^t(s) = e^{-\frac{i\pi s}{2}} \int_{C_\eta^t} \frac{e^{-z}z^{s-1}dz}{1-e^{-z}}.$$

Instead of the contour C_η^t, we consider the finite ray from the point $i\eta$ to the point z^*, see figure 7, where

$$z^* = i\eta + (t-\eta)e^{i(\frac{\pi}{2}-\epsilon)} = [\eta + (t-\eta)e^{-i\epsilon}]e^{\frac{i\pi}{2}}, \qquad 0 \le \epsilon < \frac{\pi}{2}.$$

We claim that

$$e^{-\frac{i\pi s}{2}} \int_{z^*}^{it} \frac{e^{-z}z^{s-1}dz}{1-e^{-z}} = O(t^\sigma \epsilon e^{t\epsilon^3}). \tag{6.13}$$

Indeed, the assumption $\mathrm{dist}(t, 2\pi\mathbb{Z}) > \delta$ implies that there exists an $A > 0$ independent of η, ϵ such that $|1 - e^{-z}| > A$ on the contour from z^* to it. Thus, using the parametrization $z = ite^{-i\theta}$, $0 \le \theta \le \epsilon$, in the lhs of (6.13), we can prove (6.13) as follows:

$$\begin{aligned}\left| e^{-\frac{i\pi s}{2}} \int_{z^*}^{it} \frac{e^{-z}z^{s-1}dz}{1-e^{-z}} \right| &= \left| it^s \int_0^\epsilon \frac{e^{-ite^{-i\theta}}e^{-i\theta s}d\theta}{1-e^{-ite^{-i\theta}}} \right| \le t^\sigma \int_0^\epsilon \frac{e^{-t\sin\theta}e^{\theta t}d\theta}{A} \\ &= O\left(t^\sigma \int_0^\epsilon e^{t\theta^3}d\theta\right) = O(t^\sigma \epsilon e^{t\epsilon^3}).\end{aligned}$$

---

[2] The rigorous justification of some of these arguments remains open.



Figure 7 *The contour of integration of the integral in the rhs of equation (6.14).*

where we have used that

$$|\theta - \sin\theta| \le \theta^3, \qquad 0 \le \theta \le 1.$$

It follows from (6.13) that

$$e^{-\frac{i\pi s}{2}} \int_{C_\eta^t} \frac{e^{-z} z^{s-1} dz}{1-e^{-z}} = e^{-\frac{i\pi s}{2}} \lim_{\substack{\epsilon\to 0\\ \epsilon^2 t\to 0}} \int_{\eta e^{\frac{i\pi}{2}}}^{z^*} \frac{e^{-z} z^{s-1}}{1-e^{-z}} dz. \tag{6.14}$$

We next employ the following parametrization which maps z^* to $\rho = 0$:

$$z = \eta e^{\frac{i\pi}{2}} + t\left(1 - \frac{\eta}{t} - \rho\right) e^{i\left(\frac{\pi}{2}-\epsilon\right)} = \left[\eta + t\left(1 - \frac{\eta}{t} - \rho\right) e^{-i\epsilon}\right] e^{\frac{i\pi}{2}},$$

$$0 < \rho < 1 - \frac{\eta}{t}.$$

We replace in equation (6.14) e^{-z} with the expression

$$e^{-z} = e^{-ite^{-i\epsilon} + i\rho t e^{-i\epsilon} - i\eta(1-e^{-i\epsilon})} = e^{it(\rho-1)e^{-i\epsilon} + O(\eta\epsilon)}, \qquad \epsilon \to 0$$

and we also replace z in the expression z^{s-1} of equation (6.14) with

$$z = t e^{\frac{i\pi}{2}} M e^{-i\mu} \left(1 - \frac{\rho}{M} e^{-i(\epsilon-\mu)}\right), \tag{6.15}$$



Figure 8 *The triangle D_1.*

where (M,μ) are defined via the equation

$$1+\left(\frac{t}{\eta}-1\right)e^{-i\epsilon}=\frac{t}{\eta}Me^{-i\mu}, \qquad M(t,\epsilon)>0, \quad \mu(t,\epsilon)>0. \tag{6.16}$$

Then, the rhs of equation (6.14) is given by

$$t^s \lim_{\substack{\epsilon\to 0\\ \epsilon^2 t\to 0}} e^{-i\epsilon}M^{s-1}e^{-i\mu(s-1)}\int_0^{1-\frac{\eta}{t}}\frac{e^{it(\rho-1)e^{-i\epsilon}}}{1-e^{it(\rho-1)e^{-i\epsilon}}}\left(1-\frac{\rho}{M}e^{-i\psi}\right)^{s-1}d\rho, \tag{6.17a}$$

where

$$\psi(t,\epsilon)=\epsilon-\mu. \tag{6.17b}$$

The functions M and μ are given by

$$M(t,\epsilon)=1-\frac{\epsilon^2}{2}\left(\frac{\eta}{t}-\frac{\eta^2}{t^2}\right)+O\left(\epsilon^4\left(\frac{\eta}{t}-\frac{\eta^2}{t^2}\right)^2\right), \qquad \epsilon\to 0 \tag{6.18a}$$

and

$$\mu(t,\epsilon)=\epsilon-\frac{\epsilon\eta}{t}+O(\epsilon^2), \qquad \epsilon\to 0. \tag{6.18b}$$

Indeed, consider the triangle D_1 formed by the intersection of the following finite rays, see figure 8:

$$(0,1), \quad \left(1, 1+\left(\frac{t}{\eta}-1\right)e^{-i\epsilon}\right), \quad \left(0,\frac{t}{\eta}Me^{-i\mu}\right).$$

The cosine rule for the triangle D_1 implies the following identities:

$$\left(\frac{t}{\eta}M\right)^2=1+\left(\frac{t}{\eta}-1\right)^2-2\left(\frac{t}{\eta}-1\right)\cos(\pi-\epsilon)$$

$$= 1 + \left(\frac{t}{\eta} - 1\right)^2 + 2\left(\frac{t}{\eta} - 1\right)\cos\epsilon$$
$$= \left[1 + \left(\frac{t}{\eta} - 1\right)\right]^2 - 2\left(\frac{t}{\eta} - 1\right)(1 - \cos\epsilon).$$

Thus,

$$\frac{t}{\eta}M = \frac{t}{\eta}\sqrt{1 - \frac{2\left(\frac{t}{\eta} - 1\right)(1 - \cos\epsilon)}{\frac{t^2}{\eta^2}}},$$

or

$$M = \sqrt{1 - 2\left(\frac{\eta}{t} - \frac{\eta^2}{t^2}\right)(1 - \cos\epsilon)},$$

which yields (6.18a). The sine rule for the triangle D_1 implies the identity

$$\frac{\sin\mu}{\frac{t}{\eta} - 1} = \frac{\sin(\pi - \epsilon)}{\frac{t}{\eta}M}.$$

Hence,

$$\sin\mu = \frac{1 - \frac{\eta}{t}}{M}\sin\epsilon,$$

which, using the expression (6.18a) for M, yields (6.18b).

Equation (6.18b) implies that $\psi > 0$, thus, we can define (W, w) via the equation

$$1 - \frac{\rho}{M}e^{-i\psi} = 1 + \frac{\rho}{M}e^{i(\pi - \psi)} = We^{iw}, \quad W(\rho, t, \epsilon) > 0, \quad w(\rho, t, \epsilon) > 0. \tag{6.19}$$

The functions W and w are given by

$$W(\rho, t, \epsilon) = 1 - \rho + O\left(\frac{\epsilon^2\eta}{t}\right) + O\left(\frac{\epsilon^2\eta^2}{t^2}\right), \qquad \epsilon \to 0 \tag{6.20a}$$

and

$$w(\rho, t, \epsilon) = \frac{\epsilon\eta}{t}\frac{\rho}{1 - \rho}\left[1 + O\left(\frac{\epsilon^2\eta}{t}\right) + O\left(\frac{\epsilon^2\eta^2}{t^2}\right)\right], \qquad \epsilon \to 0. \tag{6.20b}$$

Indeed, consider the triangle D_2 formed by the intersection of the following finite rays, see figure 9:

$$(0, 1), \quad \left(1, 1 + \frac{\rho}{M}e^{i(\pi - \psi)}\right), \quad \left(0, We^{iw}\right).$$

The cosine rule for the triangle D_2 implies the following identities:

$$W^2 = 1 + \frac{\rho^2}{M^2} - 2\frac{\rho}{M}\cos\psi$$
$$= \left(1 - \frac{\rho}{M}\right)^2 + \frac{2\rho}{M}(1 - \cos\psi).$$



Figure 9 *The triangle D_2.*

Hence,

$$W = \left(1 - \frac{\rho}{M}\right)\sqrt{1 + \frac{2\frac{\rho}{M}}{\left(1-\frac{\rho}{M}\right)^2}(1-\cos\psi)}.$$

Thus, using the definition (6.17b) of ψ, together with equation (6.18b), we find

$$W = 1 - \frac{\rho}{M} + \frac{1}{2}\frac{\frac{\rho}{M}}{1-\frac{\rho}{M}}\left(\frac{\epsilon^2\eta^2}{t^2} + O(\epsilon^4)\right), \qquad \epsilon \to 0.$$

Substituting in this equation the expression (6.18a) for M, we find equation (6.20a).

The sine rule for the triangle D_2 implies the identity

$$\frac{\sin w}{\rho/M} = \frac{\sin\psi}{W},$$

which, using the expressions for ψ (equations (6.17b) and (6.18b)), for M (equation (6.18a)) and for W (equation (6.20a)), yields equation (6.20b).

Using the identity

$$\left(1 - \frac{\rho}{M}e^{-i\psi}\right)^{s-1} = W^{s-1}e^{i(s-1)w},$$

as well as the expressions for M, μ, W and w obtained earlier, it is possible to compute all the expressions appearing in the rhs of (6.17a):

$$\begin{aligned}
&M^{s-1} = e^{(it+\sigma-1)\ln M} = e^{-\frac{i\epsilon^2\eta}{2}}e^{O\left(\frac{\epsilon^2}{t}\right)}e^{O\left(\frac{\epsilon^2}{t^2}\right)},\\
&e^{-i(s-1)\mu} = e^{\mu t}e^{-i(\sigma-1)\mu} = e^{\epsilon t}e^{-\epsilon\eta}e^{O(\epsilon^2 t)}e^{i(\sigma-1)\left(\frac{\epsilon\eta}{t}-\epsilon\right)}e^{O(\epsilon^2)},\\
&W^{s-1} = (1-\rho)^{s-1}e^{(it+\sigma-1)\left[O\left(\frac{\epsilon^2\eta}{t}\right)+O\left(\frac{\epsilon^2\eta^2}{t^2}\right)\right]} = (1-\rho)^{s-1}e^{O(\epsilon^2\eta)}e^{O\left(\frac{\epsilon^2\eta^2}{t}\right)},\\
&e^{iw(s-1)} = e^{-tw}e^{i(\sigma-1)w} = e^{-\epsilon\eta\frac{\rho}{1-\rho}}e^{\frac{\rho}{1-\rho}\left[O(\epsilon)+O\left(\frac{\epsilon}{t}\right)\right]}.
\end{aligned}$$

Substituting the above expressions in (6.17a) and using the identities

$$e^{-ite^{-i\epsilon}} = e^{-it(1-i\epsilon+O(\epsilon^2))} = e^{-it-\epsilon t+O(\epsilon^2 t)},$$

in order to simplify the integrand of the integral in equation (6.17a), we find the equation

$$e^{-\frac{i\pi s}{2}} \int_{C_\eta^t} \frac{e^{-z} z^{s-1} dz}{1-e^{-z}} = t^s \lim_{\substack{\epsilon \to 0 \\ \epsilon^2 t \to 0}} \int_0^{1-\frac{\eta}{t}} \frac{e^{it(\rho-1)e^{-i\epsilon}} (1-\rho)^{s-1} d\rho}{1-e^{it(\rho-1)e^{-i\epsilon}}}, \qquad 0<\eta<t. \quad (6.21)$$

The transformation $\rho = 1 - \frac{u}{t}$ maps this equation to equation (6.12).

We note that a detailed analysis in the neighborhood of $\rho = 1$ shows that similar estimates are also valid near $\rho = 1$.

A The Asymptotics of $\Gamma(1-s)$ and $\chi(s)$

The asymptotic formulae given in theorems 4.1 and 4.4 involve the functions $\chi(s)$ and $\Gamma(1-s)$. In this appendix, we derive the asymptotics of these functions as $t \to \infty$.

We use the following well known formula for the asymptotics of the Gamma function (see for example Olver [6], page 294):

$$\Gamma(s) = e^{-s} e^{s \ln s} \left(\frac{2\pi}{s}\right)^{\frac{1}{2}} \left[1 + \frac{1}{12s} + O\left(\frac{1}{s^2}\right)\right], \qquad s \to \infty, \quad |\arg s| \le \pi - \delta, \quad \delta > 0.$$

Hence, as $t \to \infty$,

$$\begin{aligned}
\Gamma(s) &= \sqrt{2\pi} e^{-\sigma - it} e^{s \ln[(it)\left(1+\frac{\sigma}{it}\right)]} \frac{1 + \frac{1}{12it\left(1+\frac{\sigma}{it}\right)} + O\left(\frac{1}{s^2}\right)}{(it)^{\frac{1}{2}} \left(1+\frac{\sigma}{it}\right)^{\frac{1}{2}}} \\
&= \frac{\sqrt{2\pi}}{\sqrt{t}} \left(e^{\frac{i\pi}{2}} t\right)^s e^{-\sigma - it} e^{(\sigma+it)\left[\frac{\sigma}{it} + \frac{\sigma^2}{2t^2} + O\left(\frac{1}{t^3}\right)\right]} e^{-\frac{i\pi}{4}} \\
&\quad \times \left[1 + \frac{1}{12it} + O\left(\frac{1}{t^2}\right)\right] \left[1 - \frac{\sigma}{2it} + O\left(\frac{1}{t^2}\right)\right] \qquad (A.1) \\
&= \sqrt{2\pi} t^{s-\frac{1}{2}} e^{\frac{i\pi s}{2}} e^{-\frac{i\pi}{4}} e^{-it} e^{-\frac{i\sigma^2}{2t} + O\left(\frac{1}{t^2}\right)} \left[1 + \frac{1-6\sigma}{12it} + O\left(\frac{1}{t^2}\right)\right] \\
&= \sqrt{2\pi} t^{s-\frac{1}{2}} e^{\frac{i\pi s}{2}} e^{-\frac{i\pi}{4}} e^{-it} \left[1 - \frac{i\sigma^2}{2t} + O\left(\frac{1}{t^2}\right)\right] \left[1 + \frac{1-6\sigma}{12it} + O\left(\frac{1}{t^2}\right)\right] \\
&= \sqrt{2\pi} t^{s-\frac{1}{2}} e^{\frac{i\pi s}{2}} e^{-\frac{i\pi}{4}} e^{-it} \left[1 + \frac{ic(\sigma)}{t} + O\left(\frac{1}{t^2}\right)\right], \qquad 0 \le \sigma \le 1, \quad t \to \infty,
\end{aligned}$$

where $c(\sigma)$ is defined by

$$c(\sigma) = \frac{\sigma}{2}(1-\sigma) - \frac{1}{12}.$$

Replacing σ by $1-\sigma$ in (A.1) and taking the complex conjugate of the resulting equation, we find

$$\Gamma(1-s) = \sqrt{2\pi} t^{\frac{1}{2}-s} e^{\frac{-i\pi(1-s)}{2}} e^{\frac{i\pi}{4}} e^{it} \left[1 - \frac{ic(\sigma)}{t} + O\left(\frac{1}{t^2}\right)\right],$$

$$0 \leq \sigma \leq 1, \quad t \to \infty. \tag{A.2}$$

On the other hand, the definition (2.2) of $\chi(s)$ implies

$$(2\pi)^s \chi(1-s) = \frac{(2\pi)^s}{2i} \frac{(2\pi)^{1-s}}{\pi} \Gamma(s) \left[e^{\frac{i\pi}{2}(1-s)} - e^{-\frac{i\pi}{2}(1-s)} \right] = \left(e^{-\frac{i\pi s}{2}} + e^{\frac{i\pi s}{2}} \right) \Gamma(s).$$

Replacing in this equation $\Gamma(s)$ by the rhs of (A.1) we find

$$\chi(1-s) = (2\pi)^{\frac{1}{2}-s} t^{s-\frac{1}{2}} e^{-\frac{i\pi}{4}} e^{-it} \left[1 + \frac{ic(\sigma)}{t} + O\left(\frac{1}{t^2}\right) \right], \qquad 0 \leq \sigma \leq 1, \quad t \to \infty. \tag{A.3}$$

Equation (A.3) together with the identity (2.15) imply the following asymptotic expression for $\chi(s)$:

$$\chi(s) = (2\pi)^{s-\frac{1}{2}} t^{\frac{1}{2}-s} e^{\frac{i\pi}{4}} e^{it} \left[1 - \frac{ic(\sigma)}{t} + O\left(\frac{1}{t^2}\right) \right], \qquad 0 \leq \sigma \leq 1, \quad t \to \infty.$$

It is of course straightforward to extend the above asymptotic formulae to higher order.

B Numerical Verifications

B.1 Verification of Theorem 3.1

Letting $\sigma = 1/2$, the error term in equation (3.2) is given by

	$t = 10$	$t = 10^2$	$t = 10^3$
$\eta = t^2$	$-(10.4 + 5.22i) \times 10^{-5}$	$(-10.2 + 2.97i) \times 10^{-9}$	$(15.1 - 4.46i) \times 10^{-13}$
$\eta = t^{\frac{3}{2}}$	$-(4.00 + 4.19i) \times 10^{-3}$	$-(1.40 + 1.11i) \times 10^{-5}$	$-(7.80 + 9.81i) \times 10^{-8}$

Note that the error is proportional to $t^3/\eta^{3+\sigma}$ as expected.

In order to demonstrate the effect of the higher order terms in (3.2), we also consider the difference between $\zeta(s)$ and the first term on the rhs of (3.2), i.e. the difference

$$\zeta(s) - \sum_{n=1}^{[\frac{\eta}{2\pi}]} n^{-s}.$$

For $\sigma = 1/2$, this difference is given by

	$t = 10$	$t = 10^2$	$t = 10^3$
$\eta = t^2$	$-0.291 + 0.274i$	$-0.341 + 0.207i$	$-0.380 + 0.121i$
$\eta = t^{\frac{3}{2}}$	$0.266 + 0.0471i$	$0.127 + 0.020i$	$0.0360 + 0.0612i$

Similarly, the difference between $\zeta(s)$ and the first two terms on the rhs of (3.2), i.e. the difference

$$\zeta(s) - \left\{ \sum_{n=1}^{[\frac{\eta}{2\pi}]} n^{-s} - \frac{1}{1-s}\left(\frac{\eta}{2\pi}\right)^{1-s} \right\},$$

is given by

	$t = 10$	$t = 10^2$	$t = 10^3$
$\eta = t^2$	$-(8.27 + 6.52i) \times 10^{-2}$	$-(6.95 + 10.3i) \times 10^{-4}$	$-(3.40 + 10.6i) \times 10^{-4}$
$\eta = t^{\frac{3}{2}}$	$(1.58 - 1.50i) \times 10^{-1}$	$(9.37 - 26.0i) \times 10^{-3}$	$(-4.93 + 3.32i) \times 10^{-3}$

The above example illustrates that the inclusion of the higher order terms in (3.2) improves the convergence of the asymptotic series considerably.

B.2 Verification of Theorem 3.2

The error term in equation (3.22) is given by

	$t = 10$	$t = 10^2$	$t = 10^3$
$\sigma = 0$	$(1.97 - 3.81i) \times 10^{-3}$	$(-7.76 + 65.1i) \times 10^{-7}$	$(5.62 - 3.40i) \times 10^{-9}$
$\sigma = 1/2$	$(2.23 - 4.34i) \times 10^{-3}$	$(-2.74 + 23.5i) \times 10^{-7}$	$(6.46 - 3.82i) \times 10^{-10}$
$\sigma = 1$	$(2.42 - 4.78i) \times 10^{-3}$	$(-9.41 + 82.5i) \times 10^{-8}$	$(7.13 - 4.22i) \times 10^{-11}$

Note that the error is proportional to $t^{-\sigma-3}$ as expected.

B.3 Verification of Theorem 4.1

Letting $\sigma = 1/2$, the error term in equation (4.2), i.e. the term

$$e^{-i\pi s}\Gamma(1-s)e^{-\frac{\pi t}{2}}\eta^{\sigma-1} \times \begin{cases} O\big(\frac{\eta}{t}\big), & 1 < \eta < t^{\frac{1}{3}} < \infty, \\ O\big(e^{-\frac{At}{\eta^2}} + \frac{\eta^4}{t^2}\big), & t^{\frac{1}{3}} < \eta < \sqrt{t} < \infty, \end{cases}$$

is given by

	$t = 10^2$	$t = 10^4$	$t = 10^6$
$\eta = 10$	$(3.04 + 7.27i) \times 10^{-3}$	$(8.05 - 2.53i) \times 10^{-6}$	$(84.1 - 5.12i) \times 10^{-10}$
$\eta = t^{1/4}$	$(45.4 - 6.75i) \times 10^{-5}$	$(8.05 - 2.53i) \times 10^{-6}$	$(-443.6 + 6.91i) \times 10^{-7}$
$\eta = t^{5/12}$	$7.27 - 1.45i) \times 10^{-1}$	$(-8.69 + 9.53i) \times 10^{-5}$	$(4.38 - 13.7i) \times 10^{-6}$

In order to illustrate the increased accuracy achieved by including the higher order terms in (4.2), we also display the corresponding table when only the first two sums on the rhs of (4.2) are included: For $\sigma = 1/2$, the difference[3]

$$\zeta(s) - \sum_{n=1}^{[\frac{t}{\eta}]} \frac{1}{n^s} - \chi(s) \sum_{n=1}^{[\frac{\eta}{2\pi}]} \frac{1}{n^{1-s}}$$

is given by

	$t = 10^2$	$t = 10^4$	$t = 10^6$
$\eta = 10$	$(5.55 - 14.1i) \times 10^{-2}$	$(-14.4 + 7.92i) \times 10^{-3}$	$(-15.9 + 4.40i) \times 10^{-4}$
$\eta = t^{1/4}$	$(4.76 - 7.53i) \times 10^{-2}$	$(-14.4 + 7.92i) \times 10^{-3}$	$-(26.3 + 6.92i) \times 10^{-3}$
$\eta = t^{5/12}$	$-(2.14 + 2.15i) \times 10^{-1}$	$(3.09 - 1.91i) \times 10^{-2}$	$(-5.70 + 10.4i) \times 10^{-3}$

B.4 Verification of Corollary 4.3

Letting $\sigma = 1/2$ and $N = 2$, we find that the error term in equation (4.22) is given by

	$t = 10$	$t = 10^2$	$t = 10^3$
$\eta = t^{\frac{7}{12}}$	$-(4.08 + 2.47i) \times 10^{-1}$	$(-2.39 + 3.50i) \times 10^{-1}$	$(-6.42 + 15.9i) \times 10^{-2}$
$\eta = t^{\frac{3}{4}}$	$(6.92 + 55.7i) \times 10^{-2}$	$(9.93 + 25.7i) \times 10^{-2}$	$-(8.75 + 12.2i) \times 10^{-3}$
$\eta = 10t$	$(1.40 - 1.12i) \times 10^{-2}$	$(2.24 + 5.28i) \times 10^{-4}$	$(2.45 + 186.1i) \times 10^{-7}$

B.5 Verification of Theorem 4.4

Letting $\sigma = 1/2$, the error term in equation (4.30), i.e. the term

$$e^{-i\pi s}\Gamma(1-s)e^{-\frac{\pi t}{2}}\eta^{\sigma-1} \times \begin{cases} O\big(\frac{\eta}{t}\big), & 1 < \eta < t^{\frac{1}{3}} < \infty, \\ O\big(e^{-\frac{At}{\eta^2}} + \frac{\eta^4}{t^2}\big), & t^{\frac{1}{3}} < \eta < \sqrt{t} < \infty, \end{cases}$$

is given by

	$t = 10^2$	$t = 10^4$	$t = 10^6$
$\eta = \sqrt{\frac{2\pi t}{100}}$	$(-5.96 + 8.83i) \times 10^{-4}$	$(6.44 + 1.30i) \times 10^{-4}$	$(4.64 + 69.7i) \times 10^{-7}$
$\eta = \sqrt{2\pi t}$	$(3.59 - 10.8i) \times 10^{-3}$	$(2.72 - 28.1i) \times 10^{-5}$	$(104.6 + 8.97i) \times 10^{-7}$
$\eta = \sqrt{200\pi t}$	$(13.7 - 5.21i) \times 10^{-6}$	$(-9.02 + 2.27i) \times 10^{-4}$	$-(14.2 + 6.74i) \times 10^{-7}$

Remark B.1 In the particular case of $\eta = \sqrt{2\pi t}$, the formula of Siegel (Eq. (32) of [8]) and the corresponding formula of Titchmarsh (Theorem 4.16 of [9]) are equivalent to the formula of our theorem 4.4. This equivalence can easily be checked numerically—for example, for $N = 3$ all three formulas yield the same numerical values for the error terms. In this regard, we note that Titchmarsh states Siegel's formula in terms of the function $\Psi(a)$ defined by

---

[3] This difference is exactly the error in the "approximate functional equation" of Hardy and Littlewood cf. [4].

$$\Psi(a) = \frac{e^{-i\pi(\frac{a^2}{2}-\frac{5}{8})}}{2\pi}\int_L \frac{e^{\frac{iw^2}{4\pi}} + aw}{e^w - 1}dw = \frac{\cos\pi(\frac{a^2}{2} - a - \frac{1}{8})}{\cos\pi a},$$

which is related to our function $\Phi(\tau, u)$ defined in (4.28) by

$$\Psi(a) = -i\Phi\Big(-1, a - \frac{1}{2}\Big)e^{-i\pi(\frac{a^2}{2}-\frac{5}{8})}, \qquad a \in \mathbb{C}.$$

Acknowledgements This work has been partially supported by the EPSRC, UK and by the Guggenheim Foundation, USA. ASF is deeply grateful to his students Michail Dimakos (partially supported by the Onassis Foundation) and Dionyssis Mantzavinos (partially supported by the EPSRC, UK) for performing a large number of numerical experiments during the early stages of this project. ASF also expresses his sincere gratitude to Anthony Ashton, Bryce McLeod, Joe Keller, David Stuart, and Eugene Shargorodsky for extensive discussions and important suggestions.

References

[1] A. C. L. Ashton and A. S. Fokas, Lindelöf's Hypothesis and the Hurwitz Zeta Function, preprint.

[2] M. V. Berry, The Riemann-Siegel expansion for the zeta function: high orders and remainders, *Proc. Roy. Soc. London Ser. A* **450** (1995), 439–462.

[3] M. V. Berry and J. P. Keating, A new asymptotic representation for $\zeta(\frac{1}{2} + it)$ and quantum spectral determinants, *Proc. Roy. Soc. London Ser. A* **437** (1992), 151–173.

[4] G. H. Hardy and J. E. Littlewood, The approximate functional equations for $\zeta(s)$ and $\zeta^2(s)$, *Proc. London Math. Soc.* **29** (1929), 81–97.

[5] J. P. Keating, In *Quantum chaos* (eds. G. Casati, I. Guarneri, & U. Smilansky), Amsterdam: North-Holland, 1993, pp. 145–185.

[6] F. W. J. Olver, *Asymptotics and Special Functions*, Academic Press, 1974.

[7] B. Riemann, Über die Anzahl der Primzahlen unter einer gegebenen Größe, Monatsberichte der Königlichen Preußischen Akademie der Wissenschaften zu Berlin (1859), 671–680.

[8] C. L. Siegel, Über Riemanns Nachlaß zur analytischen Zahlentheorie, Quellen Studien zur Geschichte der Math. Astron. und Phys. Abt. B: Studien 2: 4580 (1932), reprinted in Gesammelte Abhandlungen, Vol. 1. Berlin: Springer-Verlag, 1966.

[9] E. C. Titchmarsh, *The theory of the Riemann zeta-function*, 2nd edition, edited by D. R. Heath-Brown, The Clarendon Press, Oxford University Press, New York, 1986.